

related to the COVID-19 pandemic significantly predicted likely anxiety among Fort McMurray population. It is imperative to mobilize resources to support vulnerable communities during the COVID-19 pandemic.

Disclosure of Interest: None Declared

EPP0675

Neural Abnormalities Associated with Generalized Anxiety Disorder: A Meta-Analysis of Functional Magnetic Resonance Imaging Activation Studies

S. K. Kahlon^{1*}, Z. Ali², E. Pritchard², S. Saravia², C. Baten², A. M. Klassen², J. H. Shepherd², G. Zamora², J. Jordan¹, M. Duran², S. L. Santos², D. W. Hedges³, J. P. Hamilton⁴, M. D. Sacchet⁵ and C. H. Miller²

¹Department of Biology, Department of Psychology; ²Department of Psychology, California State University, Fresno; ³Department of Psychology, Brigham Young University, Provo, United States; ⁴Department of Biomedical and Clinical Sciences, Linköping University, Linköping, Sweden and ⁵Meditation Research Program, Department of Psychiatry, Massachusetts General Hospital, Harvard Medical School, Boston, United States

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.971

Introduction: Generalized anxiety disorder (GAD) is a highly prevalent mental illness that is associated with clinically significant distress, functional impairment, and poor emotional regulation. Primary functional magnetic resonance imaging (fMRI) studies of GAD report neural abnormalities in comparison to healthy controls. However, many of these findings in the primary literature are inconsistent, and it is unclear whether they are specific to GAD or shared transdiagnostically across related disorders.

Objectives: This meta-analysis seeks to establish the most reliable neural abnormalities observed in individuals with GAD, as reported in the primary fMRI activation literature.

Methods: We conducted an exhaustive literature search in PubMed to identify primary studies that met our pre-specified inclusion criteria and then extracted relevant data from primary, whole-brain fMRI activation studies of GAD that reported coordinates in Talairach or MNI space. We then used multilevel kernel density analysis (MKDA) with ensemble thresholding to examine the differences between adults with GAD and healthy controls in order to identify brain regions that reached statistical significance across primary studies.

Results: Patients with GAD showed statistically significant ($\alpha=0.05-0.0001$; family-wise-error-rate corrected) neural activation in various regions of the cerebral cortex and basal ganglia across a variety of experimental tasks.

Conclusions: These results inform our understanding of the neural basis of GAD and are interpreted using a frontolimbic model of anxiety as well as specific clinical symptoms of this disorder and its relation to other mood and anxiety disorders. These results also suggest possible novel targets for emerging neurostimulation therapies (e.g., transcranial magnetic stimulation) and may be used to advance our understanding of the effects of current pharmaceutical treatments and ways to improve treatment selection and symptom-targeting for patients diagnosed with GAD.

Disclosure of Interest: None Declared

EPP0676

Virtual reality exposure therapy for panic disorder

S.-E. Cho*, H. Ma and S.-G. Kang

Department of Psychiatry, Gil Medical Center, Incheon, Korea, Republic Of

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.972

Introduction: Virtual reality exposure therapy (VRET) is a treatment in a virtual environment based on the representation of a patient's disease in virtual reality. VRET means getting used to a specific situation by intentionally facing the patient's fearful situation based on the 'exposure technique.' By repeatedly exposing the patient to a fearful situation, in the end, inducing the intensity of fear to decrease without avoiding the situation. Virtual reality exposure therapy (VRET) was initially used to treat phobias, anxiety disorders, and stress. It has been proven to be an effective psychotherapy method mainly focusing on acrophobia, flight phobia, arachnophobia, social phobia, and post-traumatic stress disorder (PTSD).

Objectives: Virtual reality exposure therapy (VRET) is used to treat phobias, anxiety disorders, and stress. The virtual reality treatment system can reproduce virtual scenes with the environment modified or removed, and can familiarize patients with such environments. This study aims to verify the degree of improvement of the symptoms of panic disorder by conducting VRET that we have made considering the degree of gradual exposure to the panic disorder group and the control group.

Methods: A total of 60 subjects were included in this study, including 43 patients with panic disorder and 17 control group. Subjects were systematically exposed to specific situations over five steps. We checked the heart rate, body temperature, EEG and symptoms before and after exposure using the following assessment instruments; Subjective Units of Discomfort (SUD), Anxiety Sensitivity Index (ASI), Panic Disorder Severity Scale. Repeated measures ANOVA was performed separately for presence, SUDS (distress) scores, and PDSS across time points.

Results: In the patient group, the program proved to be effective as the sensitivity to anxiety decreased significantly after the VR program ($F=3.570$, $p<0.05$). Among the subcategories of ASI, the fear of anxiety symptoms showed statistical significance between sessions ($F=3.883$, $p<0.05$) and in the interaction between group and time ($F=4.585$, $p<0.01$). The study confirmed the effectiveness of the VR program in situations that mainly induce panic attacks in patients diagnosed with panic disorder (elevator riding, driving a car, driving a car in a tunnel, driving a car in rain). The program effect was significant through repeated measurement ANOVA for each variable of EEG (Alpha, Beta, Delta, Gamma, Theta) ($F=3.249$, $p<0.01$).

Conclusions: VRET is the process of gradually being exposed to fear stimuli, deliberately confronting and experiencing a fear situation, and learning that nothing dangerous will happen. Even if the patient experiences bodily sensations that are the subject of fear, they can reduce their fear of bodily sensations by repeatedly confirming that they are safe and getting used to them.

Disclosure of Interest: None Declared