

# ENERGY SPECTRA OF SEVERAL DISCRETE X-RAY SOURCES IN THE 20–120 keV RANGE

P. C. AGRAWAL, S. BISWAS, G. S. GOKHALE,  
V. S. IYENGAR, P. K. KUNTE, R. K. MANCHANDA and B. V. SREEKANTAN  
*Tata Institute of Fundamental Research, Bombay 5, India*

## 1. Introduction

In this paper we report on our observations of hard X-rays from several X-ray sources in the energy range 20–120 keV. The results were obtained from the data collected during two balloon flights made from Hyderabad, India (latitude  $17.6^\circ\text{N}$ , longitude  $78.5^\circ\text{E}$ ). The first flight was made on April 28, 1968, and the balloon reached a ceiling of about  $5.3\text{ g cm}^{-2}$  residual atmosphere and floated from 0230 to 0800 hrs. IST (Indian Standard Time). The second balloon was launched on December 22, 1968 and floated at about  $7.5\text{ g cm}^{-2}$  of residual air from 1000 to 1130 hrs. IST.

The X-ray telescope consisted of a NaI(Tl) crystal of area of  $97.3\text{ cm}^2$  and thickness 4 mm, coupled to a  $5''$  photomultiplier. The crystal was surrounded by both active and passive collimators. The passive collimator was a cylinder of graded shielding of lead, tin and copper and the active collimator, a cylindrical plastic scintillator. The field of view of the telescope was  $18.6^\circ$  at FWHM. The geometrical factor of the telescope for isotropic radiation was  $13.2\text{ cm}^2\text{ sr}$ . The pulses from the NaI crystal were sorted out into 10 continuous channels extending from 17 to 124 keV. An  $\text{Am}^{241}$  source came in the field of view of the telescope periodically and provided in flight calibration of the detector. All the information was recorded on a continuously moving photographic film.

The X-ray telescope was mounted on an oriented platform which was programmed to look at specified directions in the sky. The axis of the telescope was kept inclined at a fixed angle to zenith,  $25^\circ$  in the first flight and  $32^\circ$  in the second. In the first flight it was planned that the azimuth of the telescope would be aligned to the north and the south directions alternately for about 10 min each. Although the telescope looked at the directions close to north and south for a considerable period of time, it also scanned some other directions of the sky for significant period due to oscillations and 'hunting' of orienting system. Fortuitously this enabled us to make interesting observations on the X-ray intensities from these directions. A continuous record of the aspect of the telescope was made from the output of a pair of flux gate magnetometers.

In the second flight the oriented platform was programmed to look at four specified directions successively spending about 4 min in each direction during a cycle of about 16 min. The four specified directions were, N ( $\phi=0^\circ$ ), S ( $\phi=180^\circ$ ), NE ( $\phi=310^\circ$ , with the convention  $\phi=270^\circ$  being due east) and SW ( $\phi=110^\circ$ ). The performance of the orientor was very good in this flight and the telescope looked at the preselected directions of the sky successively for five cycles, from 0950 to 1115 hrs. IST.

## 2. Sco X-1

On April 28, 1968 the balloon reached ceiling altitude of  $5.3 \text{ g cm}^{-2}$  at 0230 hrs. IST. During the period 0230–0330 hrs. IST Sco X-1 was in the field of view of the telescope intermittently for a total time of 1455 sec. The angle between the telescope axis and Sco X-1 was computed every 15 sec by noting the corresponding aspect and taking into account the drift of the balloon in longitude. The effective exposure time corresponding to 100% efficiency was deduced from these data to be 255 sec. The background counting rate was determined from the counting rates during the period when no known sources were in the field of the telescope. The relevant data are given in




Fig. 1. X-ray intensities of Sco X-1 measured on April 28, 1968 and December 22, 1968.

Table I. It is seen that finite fluxes from Sco X-1 are available in four energy channels I to IV (16–72 keV) and upper limits can be put in two higher energy channels V and VI (72–125 keV). The fluxes shown in Table I are not corrected for the escape probability of K X-rays and also for the resolution and the efficiency of the detector. This correction is small and amounts to about 10%. This will be taken into account in the final analysis of the data of the X-ray sources.

During the flight on December 22, 1968, Sco X-1 was in the field of view of the

TABLE I  
Energy spectrum and intensity of Sco X-1

Date	Energy band (keV)	Mean energy (keV)	Excess counts	Exposure time (sec)	Equivalent exposure at 100% efficiency (sec)	Flux	
						Photons/keV cm <sup>2</sup> sec	keV/keV cm <sup>2</sup> sec
28.4.68	16.6-23.4	20.1	268 ± 80	1455	255	(1.43 ± 0.43) × 10 <sup>-1</sup>	2.86 ± 0.85
	23.4-31.5	27.5	72 ± 40	600	114	(1.00 ± 0.55) × 10 <sup>-2</sup>	(2.76 ± 1.6) × 10 <sup>-1</sup>
	31.5-47.7	39.6	89 ± 97	1455	255	(9.7 ± 10.6) × 10 <sup>-4</sup>	(3.85 ± 4.2) × 10 <sup>-2</sup>
	47.7-72.0	59.8	116 ± 135	1455	255	(6.0 ± 7.0) × 10 <sup>-4</sup>	(3.61 ± 4.2) × 10 <sup>-2</sup>
	72.0-97.4	84.6	2 σ = 205	1455	255	< 8.9 × 10 <sup>-4</sup>	< 7.5 × 10 <sup>-2</sup>
	97.4-123.7	110.5	2 σ = 175	1455	255	< 6.0 × 10 <sup>-4</sup>	< 6.6 × 10 <sup>-2</sup>
22.12.68	29.9-52.3	41.1	178 ± 60	566	450	(1.54 ± 0.52) × 10 <sup>-3</sup>	(6.3 ± 2.1) × 10 <sup>-2</sup>
	52.3-74.7	63.5	137 ± 79	566	450	(7.0 ± 4.0) × 10 <sup>-4</sup>	(4.5 ± 2.5) × 10 <sup>-2</sup>

telescope periodically (for about 4 min at a time) over a time period of about 90 min, from about 1000 hrs. to 1130 hrs. IST. The detector looked at Sco X-1 for a total time of 566 sec. Since the orientor performed very satisfactory the mean efficiency was as high as  $\sim 80\%$  and the effective exposure time with  $100\%$  efficiency was 450 sec. This long exposure enabled us to obtain finite and statistically significant flux in the two energy intervals of 30–52 keV and 52–75 keV (Table I), in spite of the fact that the atmospheric thickness along the line of sight to Sco X-1 was as high as  $9.2 \text{ g cm}^{-2}$  of air. In estimating the excess counts the background rate was obtained from the north region of sky which included no discrete X-ray sources.

The measured intensities of Sco X-1 during the two flights are plotted in Figure 1. In the December 22 flight rapid variations in the intensity of Sco X-1 within the period of observation of 90 min have been noticed and these results have been reported earlier [1]. However for a period of about 1 hour, the flux of Sco X-1 in the energy range 30–52 keV was fairly constant and near to the values reported by others, as well as to our flux measurements of April 28. In the figure we have plotted the flux values corresponding to this period of observation on December 22 for the two energy channels 30–52 keV and 52–75 keV.

It is seen from the Figure 1, that in the range 15 to 45 keV, the spectrum of Sco X-1




Fig. 2. X-ray intensities of Sco X-1 measured in the present experiments on 28.4.1968 and on 22.12.1968 together with available spectral information on Sco X-1.

can be represented by an exponential of the form  $N(E) = (K/E) \exp - E/E_0$ , where  $E_0 \sim 5$  keV the corresponding temperature being  $T = 5.8 \times 10^7$  K. In the energy interval 45–75 keV, our data indicate a rather flat spectrum  $E_0 \sim 20$  keV or more, suggesting that the same spectral exponent is not valid for energies below and above 45 keV.

A comparison of the present data with the available spectral data [2–8] on Sco X-1 is shown in Figure 2. It is clear from the figure that at any given energy the agreement in the absolute flux values is not better than by a factor of 2–3, at energies below 40 keV. At higher energies the spectral information on Sco X-1 is still rather poor. Our flux values at 60 and 63 keV obtained in two different flights agree with others. But the values are considerably higher than those of Buselli *et al.* [6], who also find a flattening of the spectrum at high energies ( $KT \sim 15$  keV).

### 3. Cyg X-1

On December 22, the X-ray telescope looked at Cyg X-1 in the eastern sky during the 3rd, 4th and 5th cycles. The total time of exposure at 100% efficiency was 202 sec. Excess flux was detectable in three energy channels. The background was taken corresponding to the north direction as was done in the case of Sco X-1. The flux values of Cyg X-1 are shown in Table II and plotted in Figure 3. For comparison available spectral data [8–13] are also shown in the same figure. The flux value measured in this flight at 41 keV is within the range of previously measured values. At 64 and 97 keV the present intensities appear to be higher than those measured by some investigators [11, 12, 14] but nearer to those of Overbeck and Tananbaum [8]. The spread in the flux values may be due to time variations in the intensity of Cyg X-1 also as pointed out by Overbeck and Tananbaum [8].

### 4. A New Source at High Galactic Latitude (TWX)

In April 28 flight, we had indication of some excess flux from the direction of azimuth  $90^\circ$ – $120^\circ$  and zenith angle  $25^\circ$  during the time 0210 to 0300 hrs. IST. This was clearly noticeable in the energy channel 17 to 48 keV and weakly in the energy channel 48–97.5 keV. Although the total exposure time was small (60 sec) and consequently statistical errors large, it was significant that the calculated minimum intensities of this source (assuming to be along the axis of the telescope) was almost the same as that of Sco X-1 for equivalent energy intervals. The observations in the azimuth  $90^\circ$ – $120^\circ$ , and of Sco X-1 in the azimuth  $150^\circ$ – $210^\circ$  were made intermittently over the same period of time and identical background rates were applied for both. An analysis of the data for the period 0300 to 0330 hrs. IST for the same azimuth and zenith direction, however did not show excess counts over the background level suggesting that the source probably moved out of the field of view of the telescope.

We scanned this region of the sky in the direction ( $SE(\phi = 110^\circ)$ ), on December 22, 1968 and detected considerable excess counts in the energy channels 30–52 keV and

TABLE II  
Spectrum of Cyg X-1

Date	Energy band (keV)	Mean energy (keV)	Excess counts	Exposure time (sec)	Equivalent exposure at 100% efficiency (sec)	Flux photons/keV cm <sup>2</sup> sec
December 22, 1968	29.9–52.3	41.1	149 ± 80	688	202	$(4.7 \pm 2.6) \times 10^{-3}$
	52.3–74.7	63.5	195 ± 88	688	202	$(3.4 \pm 1.6) \times 10^{-3}$
	74.7–118.7	95.7	403 ± 130	688	202	$(2.6 \pm 0.8) \times 10^{-3}$




Fig. 3. Flux values of Cyg X-1 measured on 22.12.1968 together with available flux data of Cyg X-1.

52–86 keV. The exposure time on this source region was 755 sec. During the same rotational cycles of the telescope Sco X-1 was detected in the direction S ( $\phi = 180^\circ$ ) and Cyg X-1 in the direction NW ( $\phi = 310^\circ$ ). Figure 4 shows the counting rates for the four directions scanned. Using the same value of the background counting rate as used for Sco X-1 and Cyg X-1, the flux values of the new source were calculated assuming it to be along the axis of the telescope. The flux values are given in Table III. The intensities measured on April 28 and December 22, are plotted in Figure 5 together with measurements on Sco X-1 and Cyg X-1 during these flights. The energy spectrum of the source seems to be similar to that of Sco X-1.

From the observations on April 28, the source location was deduced to be in the celestial region, R.A. = 14.0 hrs. to 15.6 hrs., and  $\delta = -5.5^\circ$  to  $+24.9^\circ$ . In the De-

TABLE III  
Intensity and energy spectrum of the new source  
(The source is assumed to be along the axis of the telescope)

Date	Energy band (keV)	Mean energy (keV)	Source counting rate-c/sec	Background counting rate-c/sec	Excess counts due to source	Total exposure Time (sec)	Flux photons/keV cm <sup>2</sup> sec
April 28, 1968	16.6–47.7	32.2	8.08	7.39	40 ± 23	60	$(2.6 \pm 1.5) \times 10^{-3}$
	47.7–97.4	72.5	13.30	12.61	38 ± 28	60	$(5.3 \pm 4.0) \times 10^{-4}$
December 22, 1968	29.9–52.3	41.1	5.29	4.98	231 ± 94	755	$(1.1 \pm 0.5) \times 10^{-3}$
	52.3–74.7	63.5	6.03	5.65	290 ± 94	755	$(8.9 \pm 2.9) \times 10^{-4}$
	74.7–118.7	96.7	11.87	11.60	203 ± 135	755	$(2.4 \pm 1.6) \times 10^{-4}$






Fig. 4. Counting rates of the X-ray telescope in the energy interval 30–119 keV on 22.12.1968 for the 4 different directions of the sky. This shows the counting rate due to the new source as compared to that of the background ( $\phi = 0^\circ$ ), Sco X-1 ( $\phi = 180^\circ$ ) and Cyg X-1 ( $\phi = 310^\circ$ ).




Fig. 5. Comparison of the spectra of the new source (TWX) and Sco X-1 measured on 28.4.1968 and 22.12.1968 and of Cyg X-1 on 22.12.1968.

ember 22 flight, the location of the source was in region, R.A. = 13.5–15.2 hrs. and  $\delta = -9.5^\circ$  to  $+23.8^\circ$ . Fortunately since the measured intensities in the two flights were nearly the same, it is estimated that the source is located in the overlapping region of the sky at R.A. = 14.0–15.2 hrs. and  $\delta = -5.5$  to  $+23.8^\circ$ , which lie in the constellations of Virgo-Bootes-Serpens. In galactic coordinates the source lies at high galactic latitude, at  $l^{\text{II}} \approx 359^\circ$  and  $b^{\text{II}} \approx +58^\circ$ .

### References

- [1] Agrawal, P. C., Biswas, S., Gokhale, G. S., Iyengar, V. S., Kunte, P. K., Manchanda, R. K., and Sreekantan, B. V.: 1969, *Nature* **224**, 51.
- [2] Chodil, G., Jopson, R. C., Mark, H., Seward, F. D., and Swift, C. D.: 1965, *Phys. Rev. Letters* **15**, 605.
- [3] Peterson, L. E. and Jacobson, A. S.: 1966, *Astrophys. J.* **145**, 62.
- [4] Lewin, W. H. G., Clark, G. W., and Smith, W. B.: 1967, *Astrophys. J. (Letters)* **150**, L153.
- [5] Chodil, G., Mark, H., Rodrigues, R., Seward, F. D., Swift, C. D., Turiel, L., Hiltner, W. A., Wallerstein, G., and Mannery, E. J.: 1968, *Astrophys. J.* **154**, 645.
- [6] Buselli, G., Clancy, M. C., Davison, P. J. N., Edwards, P. J., McCracken, K. G., and Thomas, R. M.: 1968, *Nature* **219**, 1124.
- [7] Bradt, H., Naranan, S., Rappaport, S., and Spada, G.: 1968, *Astrophys. J.* **152**, 1005.
- [8] Overbeck, J. W. and Tananbaum, H. D.: 1968, *Astrophys. J.* **153**, 899.
- [9] Grader, R. J., Hill, R. W., Seward, F. D., and Toor, A.: 1966, *Science* **152**, 1499.
- [10] Clark, G. W., Lewin, W. H. G., and Smith, W. B.: 1968, *Astrophys. J.* **151**, 21.
- [11] Peterson, L. E., Jacobson, A. S., Pelling, R. M., and Schwartz, D. A.: 1968, *Can. J. Phys.* **46**, S437.
- [12] Bleeker, J. A. M., Burger, J. J., Deerenberg, A. J. M., Scheepmaker, A., Swanenburg, B. N., and Tanaka, Y.: 1967, *Astrophys. J.* **147**, 391.
- [13] Rocchia, R., Rothenflug, R., Boclet, D., and Durochoux, Ph.: 1969, *Astron. Astrophys.* **1**, 48.