




## Research Article

# Alpha-Particle Generation from H-<sup>11</sup>B Fusion Initiated by Laser-Accelerated Boron Ions

Defeng Kong <sup>1</sup>, Shirui Xu,<sup>1</sup> Yinren Shou,<sup>1</sup> Ying Gao,<sup>1</sup> Zhusong Mei,<sup>1</sup> Zhuo Pan,<sup>1</sup> Zhipeng Liu,<sup>1</sup> Zhengxuan Cao,<sup>1</sup> Yulan Liang,<sup>1</sup> Ziyang Peng,<sup>1</sup> Pengjie Wang,<sup>1</sup> Di Luo,<sup>2,3</sup> Yang Li,<sup>2,3</sup> Zhi Li,<sup>2,3</sup> Huasheng Xie,<sup>2,3</sup> Guoqiang Zhang,<sup>4</sup> Wen Luo,<sup>5</sup> Jiarui Zhao <sup>1</sup>, Shiyong Chen,<sup>1</sup> Yixing Geng,<sup>1</sup> Yanying Zhao,<sup>1</sup> Jianming Xue,<sup>1</sup> Xueqing Yan,<sup>1,6,7</sup> and Wenjun Ma <sup>1,6,7</sup>

<sup>1</sup>State Key Laboratory of Nuclear Physics and Technology, and Key Laboratory of HEDP of the Ministry of Education, CAPT, Peking University, Beijing 100871, China

<sup>2</sup>Hebei Key Laboratory of Compact Fusion, Langfang 065001, China

<sup>3</sup>ENN Science and Technology Development Co., Ltd., Langfang 065001, China

<sup>4</sup>Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800, China

<sup>5</sup>School of Nuclear Science and Technology, University of South China, Hengyang 421001, China

<sup>6</sup>Beijing Laser Acceleration Innovation Center, Huairou, Beijing 101400, China

<sup>7</sup>Institute of Guangdong Laser Plasma Technology, Baiyun, Guangzhou 510540, China

Correspondence should be addressed to Jiarui Zhao; jrzhao@pku.edu.cn and Wenjun Ma; wenjun.ma@pku.edu.cn

Received 11 April 2022; Accepted 28 June 2022; Published 26 September 2022

Academic Editor: Fabio Belloni

Copyright © 2022 Defeng Kong et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Here, we report the generation of MeV alpha-particles from H-<sup>11</sup>B fusion initiated by laser-accelerated boron ions. Boron ions with maximum energy of 6 MeV and fluence of 10<sup>9</sup>/MeV/sr@5 MeV were generated from 60 nm-thick self-supporting boron nanofoils irradiated by 1 J femtosecond pulses at an intensity of 10<sup>19</sup> W/cm<sup>2</sup>. By bombarding secondary hydrogenous targets with the boron ions, 3 × 10<sup>5</sup>/sr alpha-particles from H-<sup>11</sup>B fusion were registered, which is consistent with the theoretical yield calculated from the measured boron energy spectra. Our results demonstrated an alternative way toward ultrashort MeV alpha-particle sources employing compact femtosecond lasers. The ion acceleration and product measurement scheme are referential for the studies on the ion stopping power and cross section of the H-<sup>11</sup>B reaction in solid or plasma.

## 1. Introduction

Nuclear fusion between proton (H) and boron (<sup>11</sup>B),  ${}^1_1\text{H} + {}^{11}_5\text{B} \rightarrow 3\alpha + 8.68 \text{ MeV}$ , is a widely concerned reaction [1–7] due to its appealing potential in fusion energy harness [8–10]. Unlike the D-D reaction and the D-<sup>6</sup>Li/D-T cycle [11], the H-<sup>11</sup>B reaction releases alpha-particles instead of neutrons, which offers clean energy without neutron radiation hazards. More importantly, <sup>11</sup>B is stable and abundant on Earth, which sheds off the fuel problem in D-T fusion. With the rapid development of high-power lasers, laser fusion based on the H-<sup>11</sup>B reaction attracts more and more

attention. However, self-sustained H-<sup>11</sup>B fusion under equilibrium conditions is highly challenging due to the insurmountable radiation loss problem at elevated temperatures. Many explorations on the H-<sup>11</sup>B reaction have been ongoing, such as driving the fusion out of thermal equilibrium by using ultrashort lasers [12–14] to reduce the radiation loss or revisiting the fusion reactivity [15–17] in the plasma environment.

In addition to the potential for clean fusion energy, the alpha-particle generation from the H-<sup>11</sup>B reaction could be a valuable source for medical and industrial applications [18–20]. The cross section for the H-<sup>11</sup>B reaction is very

large, e.g., 1.2 barn [21] at 620 keV (center-of-mass energy), and one reaction can release 1 MeV and two of 4 MeV alpha-particles [15, 22] in a simplified view. With high-energy reactants, the yield and the kinetic energies of the alpha-particles could be prominent, depending on the reaction channels. The alpha-particle generation from laser-driven H-<sup>11</sup>B reaction was firstly reported in 2005 with a yield of 10<sup>3</sup>/sr/shot [23] using a boron-rich polyethylene target irradiated by a picosecond laser. In subsequent experiments, the yields have been continuously increased to 10<sup>6</sup> α/sr/shot and 10<sup>9</sup> α/sr/shot [12, 24] in the so-called “pitcher-catcher” scheme, where energetic protons are produced from a μm-thick target through target normal sheath acceleration (TNSA) and bombard a secondary boron target.

Besides the boron-rich polyethylene targets, “sandwich” targets (SiH/B/Si) and thick boron-nitride (BN) targets were irradiated with kilojoule-scale sub-ns lasers, producing 10<sup>9</sup> and 10<sup>10</sup> /sr/shot alpha-particles, respectively [25–27]. It was found that, in spite of the difference in the driving lasers, the observed yield of the alpha-particles had a similar scaling law of about 10<sup>5</sup>–10<sup>6</sup> α/sr/.

Up to now, all the reported alpha-particle generation was driven by low-repetition rate, high-energy, long-pulse lasers. Operating one-shot typically takes an hour or more, which severely limits potential applications. Routes that employ femtosecond lasers as the drivers are noteworthy to study, which can operate at a much higher repetition rate. Besides the high-repetition rate, another advantage of femtosecond lasers is that their intensities are much higher than long-pulse lasers for the given pulse energy. The 100s TW or PW femtosecond lasers can deliver intensities of 10<sup>18</sup>–10<sup>22</sup> W/cm<sup>2</sup> on the targets. Laser-ion acceleration at such high intensity can produce copious MeV ions from nonequilibrium laser-plasma interaction, matching the cross section’s apex nicely.

Moreover, all the reported studies of laser-ion-initiated H-<sup>11</sup>B fusion utilize protons to bombard boron targets [12, 23–28]. If the opposite scheme, i.e., initiating H-<sup>11</sup>B fusion with energetic boron ions, is adopted, the generated alpha-particles would be more directional due to the higher mass of boron atoms [29]. The yield may also be enhanced as studies show that the energy conversion efficiencies from laser energy to heavy ions are higher than that of protons in favorable acceleration regimes [30, 31]. Furthermore, this scheme can be employed to investigate the stopping power of boron ions inside solid or plasma targets, which is very important for future H-<sup>11</sup>B nuclear reactors [29, 32, 33]. However, the alpha-particle generation by bombarding hydrogenous solid or plasma targets with laser-accelerated boron ions has not been realized yet. The main reason is the shortage of energetic laser-accelerated boron ions. In the widely adopted TNSA regime, the targets are μm-thick solid foils [34]. Ions with the highest charge-to-mass ratio, i.e., protons, favorably gain energy from the sheath field. The acceleration of heavy ions is drastically suppressed. With the development of laser and target-fabrication technology, ultrathin targets with nm-scale thickness were allowed to be used in the experiments, indicating the prominent efficiency for heavy ion acceleration. The variation of the laser and

target parameters leads to different regimes such as radiation pressure acceleration (RPA) [35, 36], relativistic induced transparency (RIT) [37–39], breakout afterburner acceleration (BOA) [40, 41], or hybrid acceleration [31, 42]. So far, energetic heavy ions such as C<sup>6+</sup>, Al<sup>13+</sup>, and Au<sup>51+</sup> have been produced with maximum energy up to 1.2 GeV [31, 39, 43].

In this work, we report the first H-<sup>11</sup>B fusion and alpha-particle generation results by bombarding hydrogenous targets with laser-accelerated boron ions. The MeV-level boron ions were accelerated from 60 nm-thick boron targets under the irradiation of high-contrast femtosecond laser pulses. The alpha-particles from H-<sup>11</sup>B fusion were measured by CR39 ion track detectors. The fusion reactions happening inside the hydrogenous targets are discussed considering the ion-nuclear collision, and the theoretical yield is calculated based on the measured <sup>11</sup>B spectra, which is consistent with our experimental results.

## 2. Experimental Setup

**2.1. Laser Parameters.** The experiment was performed on a 200 TW CLAPA Ti: sapphire laser system at Peking University [44]. The experimental layout is shown in Figure 1(a). An s-polarized laser pulse was normally focused on the 60 nm-thick boron nanofoils with the spot size of 8.4 × 9.2 μm (full width at half maximum) by an *f*/3 off-axis-parabolic mirror. The central wavelength and duration of the laser pulse were 800 nm and 30 fs, respectively. A cross-polarized wave system and a single plasma mirror system were employed to improve the laser contrast ratio up to 10<sup>9</sup>@40 ps and prevent the damage of targets from prepulses. The on-target laser energy was 1 J, corresponding to a peak intensity of 1 × 10<sup>19</sup> W/cm<sup>2</sup>. A 5 μm-thick plastic (C<sub>10</sub>H<sub>8</sub>O<sub>4</sub>) foil with the proton density of 4 × 10<sup>22</sup>/cm<sup>3</sup> was located 0.5 mm behind the targets at the laser axis as the “catcher” for H-<sup>11</sup>B reactions.

**2.2. Boron Nanofoil Target.** The boron nanofoils are prepared by the RF-magnetron sputtering deposition using the natural boron. The atomic ratio of <sup>10</sup>B and <sup>11</sup>B is 1 : 4. The details of the target-fabrication method will be reported elsewhere. To optimize the ion acceleration, we used 60 nm-thick self-supporting B foils, the thinnest that could be fabricated at that time, as the targets in the experimental campaign. Figure 1(b) depicts the top-view morphology of a 60 nm-thick self-supporting boron nanofoils on a target hole with a diameter of 0.5 mm. The chemical composition of the targets is characterized by an energy dispersive spectrometer in a scanning electron microscope (Figure 1(c)). Due to the oxidation of the targets in the air, the atomic ratio of B : O is 1.1 : 1. Besides, the Si atoms are from the silicon wafer as a target substrate during the fabrication. The C atoms come from the contaminated layer of nanofoils. The density of the foils, measured by the weighting method, is about 0.95 g/cm<sup>3</sup>. If the target is fully ionized, the electron density would be  $n = 160n_c$  and here the critical density would be  $n_c = m_e \omega^2 / (4\pi e^2) = 1.7 \times 10^{21} \text{ cm}^{-3}$ .




FIGURE 1: (a) Experimental setup. The laser pulses irradiate a boron nanofoil with a normal incidence. A  $5\ \mu\text{m}$ -thick plastic foil was located  $0.5\ \text{mm}$  behind the boron nanofoil to initiate the  $\text{H-}^{11}\text{B}$  fusion. The CR39, TPS, and Teflon plate were placed around the target to measure the alpha-particles and boron ions and collect the transmitted light, respectively, (b) the top-view morphology of self-supporting nanofoils three hours after preparation, and (c) the atomic weight ratio of the boron nanofoil.

**2.3. Diagnostics.** The energy spectra of the ions were measured by a Thomson parabola spectrometer (TPS) with a microchannel plate (MCP) equipped with a phosphor screen positioned  $0.78\ \text{m}$  away from the targets in the normal direction of the targets. The collimated ions with different energy and charge-to-mass ratio (CMR) were deflected by the electromagnetic fields and hit on the MCP with parabola traces. Ion signal multiplied by the MCP was converted to optical signals captured by a 16-bit EMCCD camera. For a good resolution of the traces, a tiny collimating aperture was employed. The corresponding acceptance angle is only  $4.2 \times 10^{-8}\ \text{sr}$ , which allows the recognition of single-ion events on the MCP [43, 45]. A Teflon plate with a through-hole was placed behind the target to collect the transmitted light, which can be used as a diagnostic for the laser-plasma interaction.

The alpha-particles generated from  $\text{H-}^{11}\text{B}$  fusion were detected by CR39 ion track detectors at angles of  $-45^\circ$ ,  $0^\circ$ ,  $45^\circ$ , and  $125^\circ$ . Here,  $0^\circ$  is the laser-axis direction. The distance between CR39 and the targets was  $130\ \text{mm}$ . The CR39 sheets were wrapped in  $10\ \mu\text{m}$ -thick aluminium foils to block low-energy ions. According to the Monte Carlo simulation results from SRIM [46], the minimum energy required to penetrate  $10\ \mu\text{m}$  aluminium for proton, alpha-particle, boron, carbon, and oxygen ions is  $0.8\ \text{MeV}$ ,  $2.9\ \text{MeV}$ ,  $9.5\ \text{MeV}$ ,  $12\ \text{MeV}$ , and  $16.5\ \text{MeV}$ , respectively. In our experiments, all the carbon and oxygen ions were blocked by the Al foils (see below), and only protons and alpha-particles with energy above  $0.8\ \text{MeV}$  and  $2.9\ \text{MeV}$  could go through and result in visible traces in CR39 after etching.

### 3. Result

**3.1. Energy Spectra of Borons and Other Ions.** The absolute energy spectra of boron ions can be obtained from our TPS. Figure 2(a) shows a raw image recorded by the TPS after the shooting (without the secondary plastic foil). More than ten spectral lines from boron, carbon, oxygen ions, and protons can be identified. The parabolic traces of  $^{11}\text{B}$  ions are marked with different lines. The boron ions with high charge states ( $^{11}\text{B}^{3+}$ ,  $^{11}\text{B}^{4+}$ ,  $^{11}\text{B}^{5+}$ ) can be clearly identified. Different from protons and carbon ions, the traces of boron ions are composed of cluster signals with similar shapes and clear boundaries. Due to the small acceptance angle of the TPS, the boron ions are sparsely distributed on the parabolic

traces, and a distinct cluster signal is the response of a single boron ion hitting in MCP, indicating a “single-ion” event. By summing up the counts for distinct clusters as the function of ion energy, we can obtain the response of a single boron ion [43]. Based on the single-ion response data, the absolute energy spectra of  $^{11}\text{B}^{5+}$ ,  $^{11}\text{B}^{4+}$ , and  $^{11}\text{B}^{3+}$  ions have been derived in Figure 2(b). The vertical error bars come from deviations of the single-ion response, and horizontal error bars reflect the width of the energy bins, which was adopted to  $0.2\ \text{MeV}$  to obtain smooth spectra curves. We can find that the maximum energy of  $^{11}\text{B}^{3+}$ ,  $^{11}\text{B}^{4+}$ , and  $^{11}\text{B}^{5+}$  is  $2.7\ \text{MeV}$ ,  $4.2\ \text{MeV}$ , and  $5.8\ \text{MeV}$ , respectively. The corresponding ion temperature is  $0.25\ \text{MeV}$ ,  $0.25\ \text{MeV}$ , and  $0.47\ \text{MeV}$ , respectively. The typical fluence is  $10^8$ – $10^{10}/\text{MeV}/\text{sr}$ , depending on the energy. For instance, the fluence of  $^{11}\text{B}^{5+}$  is  $10^9/\text{MeV}/\text{sr}$  at  $5\ \text{MeV}$ . The spectra of proton, carbon, and oxygen ions from this shot are given in Figure 2(c).

**3.2. Alpha-Particle Measurement.** The CR39 sheets used for alpha-particle measurement were etched in  $6\ \text{mol/L}$  NaOH solution at  $98^\circ\text{C}$  for 2 hours to reveal the ion tracks. Figure 3(a) displays the CR39 images with a solid angle of  $3.3 \times 10^{-5}\ \text{sr}$  at the angle of  $0^\circ$  and  $125^\circ$  after three shots in a row. A control CR39 sheet that was not put inside the chamber was also etched with the same procedure, whose surface morphology is shown in Figure 3(a) as well.

According to Figures 2(b) and 2(c), the maximum energy of laser-accelerated boron, carbon, and oxygen ions is  $6\ \text{MeV}$ ,  $7\ \text{MeV}$ , and  $8\ \text{MeV}$ , respectively. Therefore, those ions were completely blocked by the Al foils. The tracks of protons and alpha-particles can be easily distinguished from each other based on their sizes. We referred to the calibration of protons and alpha-particles from Zhang et al.’s works under the same etch condition [47], shown as the lines in Figure 3(b). Therefore, the dense grey dots with diameters of  $4$ – $6\ \mu\text{m}$  represent the protons, while the alpha-particles are larger black pits with diameters of  $20$ – $30\ \mu\text{m}$ , as shown in Figure 3(a). According to the proton’s spectrum from TPS and considering their energy loss in the plastic and Al foil, the proton tracks in the CR39s can be estimated as  $9 \times 10^9/\text{sr}$  at the  $0^\circ$ . So, about  $10^5$  protons can be observed in the CR39 image within a solid angle of  $3.3 \times 10^{-5}\ \text{sr}$ , consistent with the high number density of grey dots. We can find 31 alpha-particle tracks at the  $0^\circ$  direction and only 7 at  $125^\circ$ . The




FIGURE 2: (a) Raw TPS data from a 60 nm-thick boron nanofoil. The parabolic traces of  $^{11}\text{B}^{3+}$ ,  $^{11}\text{B}^{4+}$ , and  $^{11}\text{B}^{5+}$  ions have been marked with different lines, (b) ion spectra of  $^{11}\text{B}^{3+}$ ,  $^{11}\text{B}^{4+}$ , and  $^{11}\text{B}^{5+}$ , and (c) ion spectra of proton, carbon, and oxygen ions.

energy of the alpha-particles can be roughly estimated from the size of the tracks. The brown circles in Figure 3(b) show some representative alpha-particles from  $0^\circ$  direction. The energy range of alpha-particles is 3–5 MeV, which is consistent with the kinetic energy obtained from the fusion reactions. By counting the number of the alpha-particles, we can get the averaged angular distribution of alpha-particle flux per shot as can be seen in Figure 3(c). Due to the off-line measurement of CR39 and the limited beamtime, we did not perform more shots and, unfortunately, cannot give the shot-to-shot fluctuations. Generally speaking, the angular distribution shows a directional feature in the forward direction due to the momentum of the boron ions. The peak yield is  $3 \pm 0.2 \times 10^5/\text{sr}/\text{J}$  and the experimental uncertainty comes from the statistical error of tracks on CR39. It should be noted that the given values in Figure 3(c) are conservative as only alpha-particles with energies above 2.9 MeV can be detected after the shielding of the Al foils.

## 4. Discussion

We can theoretically calculate the yield of the alpha-particles from the measured boron spectra and compare it with that from the CR39 measurement. The number of boron-induced fusion reactions  $N_f$  can be estimated using the differential equation describing the ion-nuclear collisional process in the target nucleus [48] with a thickness of  $D$  represented as

$$dN_f = \int n\sigma(E)v_b(x)dt dN_b = n \int_0^D \sigma(E)dx \frac{dN_b}{dE^*} dE^*, \quad (1)$$

where  $n = 4 \times 10^{22}/\text{cm}^3$  is the proton density of the target nucleus,  $\sigma(E)$  is the nuclear cross section, and  $v_b, N_b$  are the velocity and number of incident ions, respectively.  $dN_f$  is the number of reactions driven by the boron ions impinging on the target with kinetic energy between  $E^*$  and  $E^* + dE^*$ .  $dN_b/dE^*$  represents the energy spectrum of the incident boron ions, as reported in Figure 2(b). The thickness  $D$  of




FIGURE 3: Alpha-particle generation from H-<sup>11</sup>B fusion measured by CR39. (a) Raw images of CR39 sheets, (b) calibrated track diameters versus the energy of protons and alpha-particles [47] and representative alpha-particles of our result, and (c) the angular dependence of alpha-particle flux. The inset shows the experimental layout of the pitcher-catcher scheme.

5 μm is close to the projected range  $R_{E=5.8\text{MeV}} = 7.8\mu\text{m}$  for boron ions with the maximum energy of 5.8 MeV. Although some high-energy boron ions can pass through the second target, the number is small, an order of magnitude lower than that of the 4 MeV-boron ions as shown in Figure 2(b). Moreover, their kinetic energy has degraded to below 2 MeV, corresponding to a pretty low fusion reactivity. Therefore, we believe that most boron ions are exhausted and stopped in the target nucleus for simplicity. Equation (1) can be further expressed in terms of the energy  $E$  of boron ions,

$$dN_f = n \int_{E^*}^0 \frac{\sigma(E)}{dE/dx} dE \frac{dN_b}{dE^*} dE^* = n \int_0^{E^*} \frac{\sigma(E)}{S(E)} dE \frac{dN_b}{dE^*} dE^*, \quad (2)$$

where  $S(E) = -dE/dx$  represents the stopping power of the target nucleus against incident boron ions. By integrating the energy  $E$ , the number of all alpha-particles generated from H-<sup>11</sup>B fusion can be expressed as

$$N_\alpha = 3N_f = 3n \int_0^{E_0} \left( \int_0^{E^*} \frac{\sigma(E)}{S(E)} dE \right) \frac{dN_b}{dE^*} dE^*. \quad (3)$$

Figure 4 depicts the curves of  $\sigma(E)$ ,  $S(E)$ , and  $dN_b/dE^*$  as the function of boron-ion energy.




FIGURE 4: Blackline: the stopping power  $S(E)$  of boron ions. Redline: the cross section  $\sigma(E)$  of H-<sup>11</sup>B fusion as a function of boron-ion energy in the lab. Brown lines: the energy spectra  $dN_b/dE^*$  of <sup>11</sup>B<sup>5+</sup>, <sup>11</sup>B<sup>4+</sup>, and <sup>11</sup>B<sup>3+</sup>. The dashed lines are the exponential fitting of energy spectra.

(C<sub>10</sub>H<sub>8</sub>O<sub>4</sub>) is simulated with SRIM [46], including the electronic and nuclear energy loss based on the cold target. The  $\sigma(E)$  of H-<sup>11</sup>B fusion is expressed according to Nevins and Swain’s results [21, 49] and polynomially fitted as given

TABLE 1: Curve fitting functions of  $\sigma(E)$ ,  $S(E)$ , and  $dN_b/dE^*$  and the number of alpha-particles  $N_\alpha$ .

Ions	$^{11}\text{B}^{5+}$	$^{11}\text{B}^{4+}$	$^{11}\text{B}^{3+}$	Total
$dN_b/dE^*$	$e^{(-E/0.47+30.8)}$	$e^{(-E/0.25+34.3)}$	$e^{(-E/0.25+29.2)}$	—
$\sigma(E)$	$0.0004934 E^5 - 0.0132 E^4 + 0.1155 E^3 - 0.3694 E^2 + 0.4366 E - 0.115$	—	—	—
$S(E)$	$0.0006076 E^5 - 0.02117 E^4 + 0.2801 E^3 - 1.767 E^2 + 5.194 E + 1.748$	—	—	—
$N_\alpha$	$1.04 \times 10^5$	$5.58 \times 10^4$	$3.78 \times 10^2$	$1.6 \times 10^5$
Ratio	64.9%	34.8%	0.3%	100%

<sup>†</sup>The units of the parameters match the axes in Figure 4.

in Table 1. The  $dN_b/dE^*$  of  $^{11}\text{B}^{5+}$ ,  $^{11}\text{B}^{4+}$ , and  $^{11}\text{B}^{3+}$  are also exponentially fitted in Table 1 according to Figure 2(b). The low-energy boron ions that were not measured by the TPS are also included by extrapolation down to 1 MeV. Table 1 gives the theoretical yield of alpha-particle from  $^{11}\text{B}^{5+}$ ,  $^{11}\text{B}^{4+}$ , and  $^{11}\text{B}^{3+}$  ions. One can find that the contribution from  $^{11}\text{B}^{5+}$  and  $^{11}\text{B}^{4+}$  is 64.9% and 34.8%, respectively. The energy of contribution from  $^{11}\text{B}^{4+}$  cannot be ignored even though their energy is lower than that of  $^{11}\text{B}^{5+}$ . The total yield of alpha-particles is  $N_\alpha \approx 1.6 \times 10^5/\text{sr}$ , which matches the experimental measurement from CR39 very well. Besides the  $^{11}\text{B}(p, \alpha)2\alpha$ , other channels such as  $^{12}\text{C}(p, \alpha)$  and  $^{16}\text{O}(p, \alpha)$  can also contribute to the alpha-particle generation. However, the cross section of these reactions is two to three orders of magnitude lower at the relevant energy [50]. Based on the measured energy spectra, the estimated total alpha-particle yield from the accelerated C, O, and H is about  $10^3/\text{sr}$ , two orders of magnitude lower than the observation.

The theoretical and measured alpha-particle yield of  $10^5\alpha/\text{sr}/\text{shot}$  with 1 J femtosecond laser pulses reaches a similar level to the case of proton-induced laser-driven fusion [12, 24, 25]. It should be noted that the reaction condition is still far from the apex of the cross section at 7.4 MeV ( $\sigma = 1.2$  barn) (see Figure 4). Further enhancement of the energy of B ions would lead to a higher yield and better collimation of the alpha-particles. Our simulation shows that the yield can be increased from  $1.04 \times 10^5$  to  $1.89 \times 10^6\alpha/\text{sr}/\text{shot}$  if the maximum  $^{11}\text{B}^{5+}$  energy and the temperature can be enhanced to 13 MeV and 1.3 MeV, respectively (total  $^{11}\text{B}^{5+}$  ion number keeps the same), which is very promising at higher laser intensities. Alternatively, further optimizing the thickness of the targets would also lead to higher ion energy and more H- $^{11}\text{B}$  reactions. Our 60 nm-thick B targets are slightly thicker than the optimum thickness of 10 nm in the RPA regime, according to  $l \sim \lambda a_0 n_c / n_e$ . Here,  $a_0 = 2.2$  is the normalized laser field. In these targets, TNSA probably is the primary acceleration mechanism. Further reducing the thicknesses of the targets would enable us to utilize more favorable acceleration regimes such as RPA or RIT.

## 5. Conclusion

In summary, we report the generation of  $3 \pm 0.2 \times 10^5/\text{sr}/\text{J}$  alpha-particles initiated by boron ions driven by a compact femtosecond laser for the first time. The yield is in good agreement with the theoretical calculation based on the measured  $^{11}\text{B}$  spectra, the stopping power of the boron ions in solid targets, and the reported cross section of H- $^{11}\text{B}$  fusion. Our results demonstrate an alternative way toward

ultrashort MeV alpha-particle sources with compact femtosecond lasers. The ion acceleration and product measurement scheme can provide a referential method for future studies on the stopping power of boron ions and the corresponding nuclear cross section of H- $^{11}\text{B}$  fusion in plasma by heating the “catcher” target into plasma. With higher laser intensities or thinner nanofoils in the future, the energies and the number of boron ions would further increase. The resulting higher-yield and directional alpha-particles at high-repetition rate could be promising for medical studies and industrial applications.

## Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

This work was supported by the following projects: NSFC Innovation Group Project (grant number 11921006) and National Grand Instrument Project (grant number 2019YFF01014402).

## References

- [1] D. C. Moreau, “Potentiality of the proton-boron fuel for controlled thermonuclear fusion,” *Nuclear Fusion*, vol. 17, no. 1, pp. 13–20, 1977.
- [2] J. M. Martínez-Val, S. Eliezer, M. Piera, and G. Velarde, “Fusion burning waves in proton-boron-11 plasmas,” *Physics Letters A*, vol. 216, no. 1–5, pp. 142–152, 1996.
- [3] H. Hora, G. H. Miley, M. Ghoranneviss, B. Malekynia, and N. Azizi, “Laser-optical path to nuclear energy without radioactivity: fusion of hydrogen-boron by nonlinear force driven plasma blocks,” *Optics Communications*, vol. 282, no. 20, pp. 4124–4126, 2009.
- [4] S. Eliezer and J. M. Martínez-Val, “Proton-boron-11 fusion reactions induced by heat-detonation burning waves,” *Laser and Particle Beams*, vol. 16, no. 4, pp. 581–598, 2009.
- [5] I. Last, S. Ron, and J. Jortner, “Aneutronic H+B11 nuclear fusion driven by Coulomb explosion of hydrogen nanodroplets,” *Physical Review A*, vol. 83, no. 4, 2011.
- [6] P. Lalouis, H. Hora, and S. Moustazis, “Optimized boron fusion with magnetic trapping by laser driven plasma block initiation at nonlinear forced driven ultrahigh acceleration,” *Laser and Particle Beams*, vol. 32, no. 3, pp. 409–411, 2014.

- [7] H. Hora, G. Korn, L. Giuffrida et al., G. H. Miley, S. Moustazis, and G. Mourou, Fusion energy using avalanche increased boron reactions for block-ignition by ultrahigh power picosecond laser pulses-ERRATUM,” *Laser and Particle Beams*, vol. 33, no. 4, p. 773, 2015.
- [8] N. Rostoker, M. W. Binderbauer, and H. J. Monkhorst, “Colliding beam fusion reactor,” *Science*, vol. 278, no. 5342, pp. 1419–1422, 1997.
- [9] N. Rostoker, M. W. Binderbauer, and H. J. Monkhorst, “Colliding beam fusion reactor,” in *Proceedings of the 12th International Conference on High-Power Particle Beams. BEAMS’98. Proceedings (Cat.No. 98EX103)*, pp. 195–202, Haifa, Israel, 1998.
- [10] H. Hora, G. H. Miley, M. Ghoranneviss, B. Malekynia, N. Azizic, and X.-T. He, “Fusion energy without radioactivity: laser ignition of solid hydrogen-boron (11) fuel,” *Energy & Environmental Science*, vol. 3, no. 4, pp. 478–485, 2010.
- [11] O. Hurricane, D. A. Callahan, D. T. Casey et al., J. D. Salmonson, P. T. Springer, and R. Tommasini, Fuel gain exceeding unity in an inertially confined fusion implosion,” *Nature*, vol. 506, no. 7488, pp. 343–348, 2014.
- [12] C. Labaune and C. Baccou, S. Depierreux, C. Goyon, G. Loisel, V. Yahia, and J. Rafelski, Fusion reactions initiated by laser-accelerated particle beams in a laser-produced plasma,” *Nature Communications*, vol. 4, no. 1, p. 2506, 2013.
- [13] C. Baccou, S. Depierreux, V. Yahia et al., “New scheme to produce aneutronic fusion reactions by laser-accelerated ions,” *Laser and Particle Beams*, vol. 33, no. 1, pp. 117–122, 2015.
- [14] H. Hora, S. Eliezer, G. H. Miley, J. Wang, Y. Xu, and N. Nissim, “Extreme laser pulses for non-thermal fusion ignition of hydrogen-boron for clean and low-cost energy,” *Laser and Particle Beams*, vol. 36, no. 3, pp. 335–340, 2018.
- [15] S. Stave, M. W. Ahmed, R. H. France et al., “Understanding the  $^{11}\text{B}(p, \alpha)\alpha$  reaction at the 0.675 MeV resonance,” *Physics Letters B*, vol. 696, no. 1-2, pp. 26–29, 2011.
- [16] M. H. Sikora and H. R. Weller, “A new evaluation of the  $^{11}\text{B}(p, \alpha)\alpha$  reaction rates,” *Journal of Fusion Energy*, vol. 35, no. 3, pp. 538–543, 2016.
- [17] S. V. Putvinski, D. D. Ryutov, and P. N. Yushmanov, “Fusion reactivity of the  $\text{pB}^{11}$  plasma revisited,” *Nuclear Fusion*, vol. 59, no. 7, 2019.
- [18] D.-K. Yoon, J.-Y. Jung, and T. S. Suh, “Application of proton boron fusion reaction to radiation therapy: a Monte Carlo simulation study,” *Applied Physics Letters*, vol. 105, no. 22, 2014.
- [19] L. Giuffrida, D. Margarone, G. A. P. Cirrone, A. Picciotto, G. Cuttone, and G. Korn, “Prompt gamma ray diagnostics and enhanced hadron-therapy using neutron-free nuclear reactions,” *AIP Advances*, vol. 6, no. 10, 2016.
- [20] G. A. P. Cirrone, L. Manti, D. Margarone et al., “First experimental proof of Proton Boron Capture Therapy (PBCT) to enhance protontherapy effectiveness,” *Scientific Reports*, vol. 8, no. 1, p. 1141, 2018.
- [21] W. M. Nevins and R. Swain, “The thermonuclear fusion rate coefficient for  $\text{p-}^{11}\text{B}$  reactions,” *Nuclear Fusion*, vol. 40, no. 4, pp. 865–872, 2000.
- [22] C. Labaune, C. Baccou, V. Yahia, C. Neuville, and J. Rafelski, “Laser-initiated primary and secondary nuclear reactions in Boron-Nitride,” *Scientific Reports*, vol. 6, no. 1, Article ID 21202, 2016.
- [23] V. S. Belyaev, A. P. Matafonov, V. I. Vinogradov et al., “Observation of neutronless fusion reactions in picosecond laser plasmas,” *Physical Review E: Statistical, Nonlinear, and Soft Matter Physics*, vol. 72, Article ID 026406, 2005.
- [24] J. Bonvalet, P. Nicolai, D. Raffestin et al., “Energetic  $\alpha$ -particle sources produced through proton-boron reactions by high-energy high-intensity laser beams,” *Physical Review*, vol. 103, no. 5-1, Article ID 053202, 2021.
- [25] A. Picciotto, D. Margarone, A. Velyhan et al., “Boron-proton nuclear-fusion enhancement induced in boron-doped silicon targets by low-contrast pulsed laser,” *Physical Review X*, vol. 4, no. 3, 2014.
- [26] D. Margarone, A. Picciotto, A. Velyhan et al., “Advanced scheme for high-yield laser driven nuclear reactions,” *Plasma Physics and Controlled Fusion*, vol. 57, no. 1, 2015.
- [27] L. Giuffrida, F. Belloni, D. Margarone et al., “High-current stream of energetic alpha particles from laser-driven proton-boron fusion,” *Physical Review E-Statistical Physics, Plasmas, Fluids, and Related Interdisciplinary Topics*, vol. 101, no. 1-1, Article ID 013204, 2020.
- [28] S. Kimura, A. Anzalone, and A. Bonasera, “Comment on “Observation of neutronless fusion reactions in picosecond laser plasmas,”” *Physical Review E-Statistical, Nonlinear and Soft Matter Physics*, vol. 79, no. 3 Pt 2, Article ID 038401, 2009.
- [29] A. F. Lifschitz, R. Farengo, and N. R. Arista, “Ionization, stopping, and thermalization of hydrogen and boron beams injected in fusion plasmas,” *Physics of Plasmas*, vol. 7, no. 7, pp. 3036–3041, 2000.
- [30] B. M. Hegelich, D. Jung, B. J. Albright et al., “Experimental demonstration of particle energy, conversion efficiency and spectral shape required for ion-based fast ignition,” *Nuclear Fusion*, vol. 51, no. 8, 2011.
- [31] W. J. Ma, I. J. Kim, J. Q. Yu et al., “Laser acceleration of highly energetic carbon ions using a double-layer target composed of slightly underdense plasma and ultrathin foil,” *Physical Review Letters*, vol. 122, no. 1, Article ID 014803, 2019.
- [32] R. L. Singleton, “Charged particle stopping power effects on ignition: some results from an exact calculation,” *Physics of Plasmas*, vol. 15, no. 5, Article ID 056302, 2008.
- [33] C. Deutsch and G. Maynard, “Ion stopping in dense plasmas: a basic physics approach,” *Matter and Radiation at Extremes*, vol. 1, no. 6, pp. 277–307, 2016.
- [34] S. C. Wilks, A. B. Langdon, T. E. Cowan et al., “Energetic proton generation in ultra-intense laser-solid interactions,” *Physics of Plasmas*, vol. 8, no. 2, pp. 542–549, 2001.
- [35] T. Esirkepov, M. Borghesi, S. V. Bulanov, G. Mourou, and T. Tajima, “Highly efficient relativistic-ion generation in the laser-piston regime,” *Physical Review Letters*, vol. 92, no. 17, Article ID 175003, 2004.
- [36] X. Q. Yan, C. Lin, Z. M. Sheng et al., “Generating high-current monoenergetic proton beams by a circularly polarized laser pulse in the phase-stable acceleration regime,” *Physical Review Letters*, vol. 100, no. 13, Article ID 135003, 2008.
- [37] A. Henig, D. Kiefer, K. Markey et al., “Enhanced laser-driven ion acceleration in the relativistic transparency regime,” *Physical Review Letters*, vol. 103, no. 4, Article ID 045002, 2009.
- [38] B. M. Hegelich, I. Pomerantz, L. Yin et al., “Laser-driven ion acceleration from relativistically transparent nanotargets,” *New Journal of Physics*, vol. 15, no. 8, Article ID 085015, 2013.
- [39] S. Palaniyappan, C. Huang, D. C. Gautier et al., “Efficient quasi-monoenergetic ion beams from laser-driven relativistic plasmas,” *Nature Communications*, vol. 6, no. 1, Article ID 10170, 2015.
- [40] L. Yin, B. J. Albright, B. M. Hegelich, and J. C. Fernández, “GeV laser ion acceleration from ultrathin targets: the laser

- break-out afterburner,” *Laser and Particle Beams*, vol. 24, no. 2, pp. 291–298, 2006.
- [41] L. Yin, B. J. Albright, K. J. Bowers, D. Jung, J. C. Fernández, and B. M. Hegelich, “Three-dimensional dynamics of breakout afterburner ion acceleration using high-contrast short-pulse laser and nanoscale targets,” *Physical Review Letters*, vol. 107, no. 4, Article ID 045003, 2011.
- [42] A. Higginson, R. J. Gray, M. King et al., “Near-100 MeV protons via a laser-driven transparency-enhanced hybrid acceleration scheme,” *Nature Communications*, vol. 9, no. 1, p. 724, 2018.
- [43] P. Wang, Z. Gong, S. G. Lee et al., “Super-heavy ions acceleration driven by ultrashort laser pulses at ultrahigh intensity,” *Physical Review X*, vol. 11, no. 2, 2021.
- [44] Y.-X. Geng, Q. Liao, Y.-R. Shou et al., “Generating proton beams exceeding 10 MeV using high contrast 60 TW laser,” *Chinese Physics Letters*, vol. 35, no. 9, 2018.
- [45] M. Nishiuchi, N. P. Dover, M. Hata et al., “Dynamics of laser-driven heavy-ion acceleration clarified by ion charge states,” *Physical Review Research*, vol. 2, no. 3, 2020.
- [46] J. F. Ziegler, M. D. Ziegler, and J. P. Biersack, “SRIM—the stopping and range of ions in matter,” *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, vol. 268, no. 11–12, pp. 1818–1823, 2010.
- [47] Y. Zhang, H.-W. Wang, Y.-G. Ma et al., “Energy calibration of a CR-39 nuclear-track detector irradiated by charged particles,” *Nuclear Science and Techniques*, vol. 30, no. 6, 2019.
- [48] K. S. Krane, *Introductory Nuclear Physics*, John Wiley & Sons, Hoboken, NJ, USA, 1991.
- [49] H. Becker, C. Rolfs, and H. Trautvetter, “Low-energy cross sections for  $^{11}\text{B}(p, 3\alpha)$ ,” *Zeitschrift für Physik A Atomic Nuclei*, vol. 327, no. 3, pp. 341–355, 1987.
- [50] A. Whitehead and J. Foster, “Activation cross sections for  $\text{C}^{12}(p, pn)\text{C}^{11}$ ,  $\text{O}^{16}(p, \alpha)\text{N}^{13}$ , and  $\text{F}^{19}(p, pn)\text{F}^{18}$ ,” *Canadian Journal of Physics*, vol. 36, no. 10, pp. 1276–1285, 1958.