



## Vitamin E–drug interactions: molecular basis and clinical relevance

Maren Podszun and Jan Frank\*

*Institute of Biological Chemistry and Nutrition, University of Hohenheim, D-70599 Stuttgart, Germany*

### Abstract

Vitamin E ( $\alpha$ -,  $\beta$ -,  $\gamma$ - and  $\delta$ -tocopherol and -tocotrienol) is an essential factor in the human diet and regularly taken as a dietary supplement by many people, who act under the assumption that it may be good for their health and can do no harm. With the publication of meta-analyses reporting increased mortality in persons taking vitamin E supplements, the safety of the micronutrient was questioned and interactions with prescription drugs were suggested as one potentially underlying mechanism. Here, we review the evidence in the scientific literature for adverse vitamin E–drug interactions and discuss the potential of each of the eight vitamin E congeners to alter the activity of drugs. In summary, there is no evidence from animal models or randomised controlled human trials to suggest that the intake of tocopherols and tocotrienols at nutritionally relevant doses may cause adverse nutrient–drug interactions. Consumption of high-dose vitamin E supplements ( $\geq 300$  mg/d), however, may lead to interactions with the drugs aspirin, warfarin, tamoxifen and cyclosporine A that may alter their activities. For the majority of drugs, however, interactions with vitamin E, even at high doses, have not been observed and are thus unlikely.

**Key words:** Drug metabolism: Interactions: Pharmacodynamics: Pharmacokinetics: Tocopherols: Tocotrienols: Vitamin E

### Introduction

Many individuals take vitamin E supplements under the assumption that they may be good for their health or, if not that, at least can do no harm. The well-documented safety of the micronutrient<sup>(1)</sup> started being questioned with the publication of meta-analyses reporting increased mortality in individuals taking high-dose vitamin E supplements<sup>(2,3)</sup> and interactions with prescription drugs were proposed as one possible explanation for these adverse events<sup>(4)</sup>. Here, we review the evidence available in the scientific literature for adverse vitamin E–drug interactions and their underlying mechanisms.

### Vitamin E


The term vitamin E refers to the eight lipid-soluble substances  $\alpha$ -,  $\beta$ -,  $\gamma$ - and  $\delta$ -tocopherol (T) and -tocotrienol ( $T_3$ ) (Fig. 1), which are exclusively produced by photosynthetic organisms and therefore present at varying concentrations in most plant foods<sup>(5)</sup>. Chemically, T and  $T_3$  consist of a chromanol ring attached to a sixteen-carbon saturated alkyl (T), or a threefold unsaturated isoprenoid

side chain ( $T_3$ ), respectively. The T side chain has three chiral centres at positions 2, 4' and 8', which can be either in the R or S conformation and make possible eight different stereoisomers (RRR, RSR, RRS, RSS, SRR, SSR, SRS and SSS).  $T_3$  possess only one chiral centre at position 2, which can be either in the R or S configuration. The biosynthesis of vitamin E in plants is stereo-selective and therefore natural T are exclusively in the RRR configuration, whereas synthetic T consist of an equimolar (all racemic; all rac) mixture of all eight possible stereoisomers.  $T_3$  are not synthesised, but extracted from natural sources, and thus present in the diet as stereoisomers with the R conformation at carbon 2 (2R). The prefixes  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  denote the number and positions of methyl groups substituted at the chromanol ring (Fig. 1)<sup>(5,6)</sup>.

More than 90 years ago, vitamin E was identified as a dietary factor required for maintaining fertility in female rats by preventing fetal resorption<sup>(7)</sup>. The anti-sterility activity of the eight vitamin E congeners differs in rats, with  $\alpha$ T being the most active<sup>(8)</sup>. Furthermore,  $\alpha$ T is the major lipid-soluble antioxidant in human plasma<sup>(9)</sup> and may (thereby) modulate gene expression and cellular

**Abbreviations:** all rac, all racemic; COX, cyclo-oxygenase; CYP, cytochrome P<sub>450</sub> mixed-function mono-oxygenase; GST, glutathione-S-transferase; NQO1, NAD(P)H:quinone oxidoreductase; OATP, organic anion transporting polypeptide; P-gp, P-glycoprotein; PKC, protein kinase C; PMA, phorbol myristate acetate; PXR, pregnane X receptor; T, tocopherol;  $T_3$ , tocotrienol; TXA2, thromboxane A2; UGT, UDP-glucuronosyltransferase.

\* **Corresponding author:** Professor Jan Frank, fax +49 711 459 23386, email jan.frank@nutres.de



**Fig. 1.** The chemical structures of tocopherols and tocotrienols.

signalling<sup>(10–12)</sup>. The biochemical mechanism underlying its essential vitamin function, however, is still unknown.

Although all congeners are ingested with the diet, the organism selectively retains  $\alpha$ T and preferentially metabolises and excretes the non- $\alpha$ T congeners<sup>(13)</sup>. Therefore,  $\alpha$ T is considered the most important form of vitamin E<sup>(14)</sup> and is the congener typically found in dietary supplements.

In Germany, 28 % of the population regularly take dietary supplements and 11 % frequently take vitamin E<sup>(15)</sup>. Dietary supplement use may be even higher in senior citizens. In a Southern German cohort of individuals aged 65 years or older, 46 % took dietary supplements and 13 % supplemented vitamin E<sup>(16)</sup>.

In the USA, almost 40 % of the population take over-the-counter vitamin or mineral supplements, of which approximately 37 % contain vitamin E<sup>(17)</sup>. These dietary supplements may contain vitamin E in doses as high as 1000 IU per serving; equivalent to 45–60 times the RDA of 18–22 IU/d for adults in the USA<sup>(14,18)</sup> or the 16–22 IU/d recommended by the German, Austrian and Swiss nutrition societies<sup>(19)</sup>.

Various meta-analyses questioned the previously established safety (in doses up to 800 IU/d) of vitamin E and suggested an increased risk of gastrointestinal cancer, heart failure, and all-cause mortality in subjects consuming dietary supplements containing vitamin E in high doses (for a review, see Bell & Grochoski<sup>(20)</sup>). Many of the intervention trials included in the meta-analyses studied patients with pre-existing morbidities who took one or more prescription drugs alongside dietary supplements (for reviews, see Bell & Grochoski<sup>(20)</sup> and Frank & Rimbach<sup>(21)</sup>). In some randomised, controlled human intervention trials, vitamin E was even administered in combination with a drug as part of the study design (for a discussion, see Brigelius-Flohe<sup>(4,22)</sup>, Bell & Grochoski<sup>(20)</sup> and Frank & Rimbach<sup>(21)</sup>). Hence, many of these human studies may have suffered from bias due to adverse effects caused by nutrient–drug interactions (for a review, see Frank & Rimbach<sup>(21)</sup>), which questions the validity of the conclusions drawn from the meta-analyses of these trials<sup>(1,20)</sup>.

### Nutrient–drug interactions


In order to reach the systemic circulation, orally ingested nutrients and drugs are required to pass a number of

physiological barriers. They first need to be liberated from the food matrix or drug formulation, mainly in the stomach and small intestine, before they can pass the second barrier, absorption into gastrointestinal cells. In the endothelial cells of the gastrointestinal tract, dietary factors and drugs may undergo metabolism by phase I and II enzymes, and both the parent compounds and their metabolites may either be excreted back into the intestinal lumen by membrane transporters on the luminal side, or secreted into the mesenteric vein or lymph. Compounds present in the mesenteric blood or lymph ultimately reach the liver via the portal vein. In the liver, the parent compounds and/or metabolites can be further metabolised by hepatic phase I and II enzymes and are either excreted with the bile back into the small intestine or are secreted into the systemic circulation, from where they are distributed through the body. If they are able to pass through the vascular endothelium, they are taken up into peripheral tissues, where they may exert their biological or pharmacological function. Compounds that are not taken up by peripheral tissues are either returned to the liver or delivered to the kidney for renal elimination.

With the exception of parenteral nutrition, nutrients and other dietary factors are generally ingested orally. A drug, however, may also be administered by injection, inhalation or other routes, which may reduce the number of barriers it has to pass before reaching the systemic circulation.

The term pharmacokinetics (‘what the body does to the drug’) is used to describe the extent (concentration) to which a drug or its active metabolite is present in the systemic circulation and how long it stays there. The term pharmacodynamics (‘what the drug does to the body’), on the other hand, designates the biological activity of a drug, for example, the lowering of blood lipids, inhibition of platelet aggregation and so forth<sup>(23)</sup>.

Interactions between a nutrient and a drug may occur in either of two ways: a nutrient may alter the pharmacokinetics and/or pharmacodynamics of a drug (nutrient–drug interaction) or a drug may alter the concentration and/or biological activity of a nutrient in the organism (drug–nutrient interaction)<sup>(24)</sup>. Drug–nutrient interactions, such as impaired absorption of vitamin E when administered together with drugs that reduce fat-absorption, are not within the scope of the present review. Nutrient–drug



**Fig. 2.** Effects of vitamin E observed in cell-culture, animal and/or human trials on targets potentially altering the pharmacokinetics and pharmacodynamics of drugs. P-gp, P-glycoprotein; CYP, cytochrome P<sub>450</sub> mixed-function mono-oxygenase; UGT, UDP-glucuronosyltransferase; T, tocopherol; ↔, no change; T<sub>3</sub>, tocotrienol; ↑, increase; OATP, organic anion transporting polypeptide; ↓, decrease. For detailed information on the experiments and outcomes, see Tables 1 and 2. (A colour version of this figure can be found online at [www.journals.cambridge.org/nrr](http://www.journals.cambridge.org/nrr)).

interactions can be roughly categorised, based on the underlying mechanism and nature of the interaction, into: (1) drug inactivation occurring *ex vivo* (in the drug formulation and outside of the body); (2) interactions affecting pre-systemic transport (uptake) and metabolism; (3) interactions affecting systemic metabolism and elimination of the drug; and (4) interactions directly altering drug activity (pharmacodynamics). However, overlap between these hypothetical classes may, and is likely to, occur. Interactions of the second and third kind alter the pharmacokinetics and interactions of the fourth type the pharmacodynamics of drugs (Fig. 2).

Nutrient–drug interactions, that is, a decrease or increase in the concentration of the active drug and/or altered pharmacodynamics, may increase the risk for and rate of adverse effects. In the present review, we summarise and discuss the evidence for potentially adverse interactions of vitamin E with drugs (nutrient–drug interactions) according to these four categories.

## Potential interactions of vitamin E with drugs

### *Ex vivo inactivation of drugs*

The first class of nutrient–drug interactions comprises chemical reactions (for example, complexation, hydrolysis, precipitation, oxidation) between a nutrient and a drug, which occur within the galenic formulation and mainly outside of the body<sup>(24)</sup>. T and T<sub>3</sub> are potent antioxidants

in homogeneous non-aqueous solutions and liposomal formulations<sup>(6)</sup> and could thus protect easily oxidisable drugs with reduction potentials higher than those of the respective vitamin E congener. For drugs with reduction potentials lower than those of vitamin E, the nutrient could act as oxidising agent and, at least theoretically, oxidatively damage the drug molecule<sup>(25)</sup>. T and their radical forms (tocopheroxyl radicals) are capable of reducing transition metal ions to a lower valence state (for example, reduction of Fe<sup>3+</sup> to Fe<sup>2+</sup>), which could favour pro-oxidant reactions catalysed by these metals<sup>(6)</sup>. Hence, drug formulations with considerable amounts of free transition metals could theoretically be at risk of increased Fenton or Fenton-like oxidation reactions. To date, however, no vitamin E–drug interactions of this kind have been reported in the scientific literature.

### *Vitamin E and pharmacokinetics*

**Pre-systemic transport and metabolism.** The second class of interactions discussed considers interactions that alter the absorption of a drug and mostly occur before the drug appears in the systemic circulation. Such interactions may be facilitated by modulation of the activity of membrane transporters or enzymes involved in xenobiotic metabolism or drug binding<sup>(24)</sup> and occur on the level of the pre-systemic transport and metabolism.

**Vitamin E and pre-systemic drug transport.** Epithelial efflux transporters, such as the multidrug resistance



**Table 1.** Effects of vitamin E on mRNA or protein expression or activity of transporters and enzymes potentially affecting the pharmacokinetics of drugs in cell-culture, animal and human experiments

Vitamin E congener	Model	Treatment and dose	Duration	Effects on transporters and enzymes			Reference
				Phase I enzymes	Phase II enzymes	Membrane transporters	
Intestine Synthetic (all rac) $\alpha$ T, $\beta$ T, $\gamma$ T, $\delta$ T, $\alpha$ T <sub>3</sub> , $\beta$ T <sub>3</sub> , $\gamma$ T <sub>3</sub> , $\delta$ T <sub>3</sub>	Human colon cancer cell line LS180	Incubation, 10 $\mu$ mol/l	24 h	mRNA expression: CYP3A4 $\leftrightarrow$ for all congeners	mRNA expression: UGT1A1 $\alpha$ T <sub>3</sub> , $\beta$ T <sub>3</sub> , $\delta$ T <sub>3</sub> $\uparrow$ (1.5- to 2.0-fold) $\alpha$ T, $\beta$ T, $\gamma$ T, $\delta$ T, $\gamma$ T <sub>3</sub> $\leftrightarrow$	mRNA expression: P-gp $\alpha$ T <sub>3</sub> , $\beta$ T <sub>3</sub> , $\gamma$ T <sub>3</sub> , $\delta$ T <sub>3</sub> $\uparrow$ (2- to 4-fold) $\alpha$ T, $\beta$ T, $\gamma$ T, $\delta$ T $\leftrightarrow$	(30)
RRR- $\gamma$ T <sub>3</sub>	Human colon cancer cell line LS180	Incubation, 25 $\mu$ mol/l	48 h			Protein expression: P-gp $\uparrow$ (2.5-fold) P-gp activity: Efflux of rhodamin 123 $\uparrow$	(31)
RRR- $\alpha$ T	Guinea-pigs	Feeding, 250 mg/kg diet	6 weeks	Protein expression: CYP3A4 $\leftrightarrow$			(37)
$\alpha$ T (stereochemistry not specified)	Human colon cancer cell line Colo205	Incubation, 0.0001–10 $\mu$ mol/l	48 h		GST, NQO1 activity: GST $\leftrightarrow$ NQO1 $\uparrow$ (dose-dependent)		(38)
$\alpha$ T (stereochemistry not specified)	Rats (microsomes of small and large intestine)	Feeding, 200 mg/kg diet	2 weeks		Total UGT activity: 4-methylumbelliferone glucuronidation $\leftrightarrow$		(40)
Liver RRR- $\alpha$ T	Rats	Subcutaneous injection, 100 mg/kg BW per d	7 d			mRNA expression: OATP C $\downarrow$	(45)
RRR- $\alpha$ T	Guinea-pigs	Feeding, 250 mg/kg diet	6 weeks			Protein expression: OATP C $\leftrightarrow$ OATP C activity: Atorvastatin plasma concentration $\leftrightarrow$	(37)
Synthetic (all rac) $\alpha$ T, $\beta$ T, $\gamma$ T, $\delta$ T, $\alpha$ T <sub>3</sub> , $\beta$ T <sub>3</sub> , $\gamma$ T <sub>3</sub> , $\delta$ T <sub>3</sub>	Human primary hepatocytes	Incubation, 10 $\mu$ mol/l	24 h	mRNA expression: CYP3A4 $\alpha$ T <sub>3</sub> , $\beta$ T <sub>3</sub> , $\gamma$ T <sub>3</sub> , $\delta$ T <sub>3</sub> $\uparrow$ (3- to 5-fold) $\alpha$ T, $\beta$ T, $\gamma$ T, $\delta$ T $\leftrightarrow$	mRNA expression: UGT1A1 $\leftrightarrow$ for all	mRNA expression: P-gp $\uparrow$ $\leftrightarrow$ for all	(30)
RRR- v. all rac- $\alpha$ T acetate	Human liver cancer cell line HepG2	Incubation, 0–300 $\mu$ mol/l	7 d (medium changed every 24 h)	mRNA expression: CYP1A2, 2A6, 2C9, 3A4, 3A5, 3A7, 3A43, 4F3, 4F12, 11B1, 24A1, 27A1, 51A1 RRR- and all rac- $\alpha$ T $\leftrightarrow$ CYP20A1 $\uparrow$ (dose-dependent up-regulation by RRR- and all rac- $\alpha$ T)			(55)
RRR- $\alpha$ T acetate	Mice	Feeding, 2 v. 20 mg/kg diet	3 months	mRNA expression: Cyp3a11* $\downarrow$			(56)
RRR- $\alpha$ T acetate	Mice	Feeding, 20 v. 200 mg/kg diet	9 months	mRNA expression: Cyp3a11* $\leftrightarrow$			(56)
RRR- $\alpha$ T acetate + $\gamma$ T <sub>3</sub>	Mice	Feeding, 20 v. 200 mg/kg diet +250 $\mu$ g/d $\gamma$ T <sub>3</sub>	7 d	mRNA expression: Cyp3a11* $\leftrightarrow$			(56)
RRR- $\alpha$ T acetate	Rats	Feeding, 1.7 v. 60 mg/kg diet	290 d	mRNA expression: CYP1A2, 2A1, 2A2, 2B2, 2B3, 2C6, 2C7, 2C11, 2C13 2C22, 2C23, 2D3, 2D4, 2D5, 2F4, 2J3, 3A1*, 3A18*, 4A1, 4F1, 4F4, 4F6, 7A1 $\leftrightarrow$			(55)
All rac- $\alpha$ T acetate	Mice	Feeding, 35 v. 1000 mg/kg diet	4 months	mRNA expression: Cyp3a11* $\uparrow$ (1.3- to 4-fold)	mRNA expression: Sult2a $\uparrow$ (10.8- to 10.9-fold) Gstm3 $\uparrow$ (1.8- to 1.9-fold)	mRNA expression: Mdr1a $\uparrow$ (1.9- to 2.0-fold)	(57)



**Table 1.** Continued

Vitamin E congener	Model	Treatment and dose	Duration	Effects on transporters and enzymes			Reference
				Phase I enzymes	Phase II enzymes	Membrane transporters	
RRR- $\alpha$ T	Guinea-pigs	20 v. 250 mg/kg diet	6 weeks	Protein expression: CYP3A4, 4F2, 20A1 $\leftrightarrow$ , CYP3A4 activity: Parent and metabolite plasma concentration of atorvastatin $\leftrightarrow$			(37)
RRR- $\alpha$ T acetate	Hypercholesterolaemic patients on lovastatin or simvastatin therapy (CYP3A4 substrates)	400 mg/d (dietary supplement)	8 weeks	CYP3A activity: 6 $\beta$ -hydroxycortisol/cortisol ratio $\leftrightarrow$			(58)
RRR- $\alpha$ T	Healthy human subjects	Supplementation of 503 mg/d	3 weeks	CYP3A activity: Blood AUC of the CYP3A4 substrate midazolam $\leftrightarrow$			(59)
$\alpha$ T (stereochemistry not specified)	Rats	Feeding, 200 mg/kg diet	2 weeks		UGT activity 4-methylumbelliferone-glucuronidation $\leftrightarrow$		(40)
$\alpha$ T (stereochemistry not specified)	Rats	Feeding, 2500 mg/kg diet	10 d		GST activity 1-Chloro-2,4-dinitrobenzene glutathionylation $\uparrow$ (1.76-fold) 2,4-Dichloro-1-nitrobenzene glutathionylation $\uparrow$ (2.68-fold)		(60)
$\alpha$ T, $\delta$ T, $\alpha$ T <sub>3</sub> , $\gamma$ T <sub>3</sub>	Isolated human glutathione-S-transferase P1-1				GST activity $\downarrow$ GST P1-1 IC <sub>50</sub> : $\alpha$ T, 0.7; $\delta$ T, 0.8; $\alpha$ T <sub>3</sub> , 1.8; $\gamma$ T <sub>3</sub> , 0.7 $\mu$ mol/l		(61,62)
$\alpha$ T, micellar formulation	Rats	Subcutaneous injections of 100 mg/kg BW; saline injections as control	9 d			mRNA expression: ABCB1 (P-gp) $\uparrow$ (10-fold)	(45)
$\alpha$ T, ethanolic solution	Rats	Subcutaneous injections of 100 mg/kg BW; saline injections as control	7 d			mRNA expression: ABCB1 (P-gp) $\uparrow$ (3.57-fold) Protein expression: ABCB1 (P-gp) $\uparrow$	(45)

All rac, all racemic; T, tocopherol; T<sub>3</sub>, tocotrienol; CYP, cytochrome P<sub>450</sub> mixed-function mono-oxygenase;  $\leftrightarrow$ , no change; UGT, UDP-glucuronosyltransferase;  $\uparrow$  increase; P-gp, P-glycoprotein; GST, glutathione-S-transferase; NQO1, NAD(P)H:quinone oxidoreductase; BW, body weight; OATP, organic anion transporting polypeptide;  $\downarrow$ , decrease; Sult2a, sulfotransferase 2A; Mdr1a, multidrug resistance protein 1a; Gstm3, glutathione S-transferase M3; IC<sub>50</sub>, 50% inhibition; ABCB1, ATP-binding cassette transporter B1.

\* Homologous to human CYP3A4.



**Table 2.** Effects of vitamin E on the activity (pharmacodynamics) of drugs in cell culture, animal and human experiments

Vitamin E congener	Model	Treatment and dose	Time	Experiment	Results	Reference
RRR- $\alpha$ T succinate	Platelets isolated from healthy human subjects	0.016–16.22 $\mu$ mol/l	Pre-incubation with $\alpha$ T for 5 min	Cyclo-oxygenase activity (arachidonic acid as substrate)	Dose-dependent reduction in cyclo-oxygenase activity: Thromboxane A <sub>2</sub> ↓ PGD <sub>2</sub> ↓	(66)
RRR- $\alpha$ T, RRR- $\alpha$ T acetate, all rac- $\alpha$ T acetate	Platelet-rich plasma isolated from healthy human subjects	500 $\mu$ mol/l	Pre-incubation with $\alpha$ T for 30 min	Platelet aggregation induced by addition of arachidonic acid or PMA	Induced aggregation: RRR- $\alpha$ T ↓ RRR- $\alpha$ T acetate ↓ All rac- $\alpha$ T acetate ↔ Protein kinase C-mediated protein phosphorylation: $\alpha$ T ↓ RRR- $\alpha$ T acetate ↓ All rac- $\alpha$ T acetate ↔	(68)
RRR- $\alpha$ T	Daily supplementation of healthy human subjects; <i>n</i> 15	267, 533 or 800 mg/d	2 weeks	<i>Ex vivo</i> aggregation of isolated platelets induced by arachidonic acid, PMA or ADP	ADP-induced aggregation: All doses ↔ Arachidonic acid-induced aggregation: 267 and 533 mg/d ↔ 800 mg/d ↓ PMA-induced aggregation: All doses ↓	(67)
All rac- $\alpha$ T v. mixed tocopherols v. untreated control	Healthy human subjects; <i>n</i> 48	100 mg/d v. 20 mg RRR- $\alpha$ T, 100 mg RRR- $\gamma$ T, and 40 mg $\delta$ T per d	8 weeks	Aggregation of isolated platelets with ADP or PMA	ADP-induced aggregation: All rac- $\alpha$ T ↔ Mixed T ↓ PMA-induced aggregation: All rac- $\alpha$ T ↔ Mixed T ↔ Activation of PKC: All rac- $\alpha$ T ↓ Mixed T ↓	(69)
All rac- $\alpha$ T acetate	Healthy human subjects; <i>n</i> 47 (twenty males and twenty-seven females)	400 mg/d for 2 weeks, 800 mg/d for 2 weeks, and 1200 mg/d for 2 weeks v. untreated control	6 weeks	<i>Ex vivo</i> adhesion of isolated platelets to collagen <i>Ex vivo</i> aggregation of isolated platelets induced by collagen, adrenaline or ADP	Platelet adhesion: Dose-dependently ↓ Platelet aggregation: Adrenaline- and ADP-induced aggregation modestly but not significantly ↓ Collagen-induced aggregation ↓ only in females	(70)
All rac- $\alpha$ T acetate	400 mg/d for 2 weeks, 800 mg/d for 2 weeks, and 1200 mg/d for 2 weeks alone v. the same treatment plus 300 mg acetylsalicylic acid every other day for the entire 6 weeks	6 weeks	Healthy human subjects; <i>n</i> 47 (twenty males and twenty-seven females)	<i>Ex vivo</i> adhesion of isolated platelets to collagen <i>Ex vivo</i> aggregation of isolated platelets induced by collagen, adrenaline or ADP	Platelet adhesion: ↔ Compared with either T or acetylsalicylic acid alone Platelet aggregation: ↔ Compared with either T or acetylsalicylic acid alone	(70)
All rac- $\alpha$ T acetate	Male smokers, <i>n</i> 409 (subset of the Alpha-Tocopherol Beta-Carotene Cancer Prevention Study)	50 mg/d	5–7 years	Supplementation with all rac- $\alpha$ T acetate in combination with 100–3200 mg/d acetylsalicylic acid	Increased oral 'bleeding on probing' with $\alpha$ T supplementation compared with acetylsalicylic acid alone	(72)
All rac- $\alpha$ T acetate	Patients who previously suffered from an ischaemic event; <i>n</i> 50	400 mg	≤ 2 years	Combination with acetylsalicylic acid (325 mg)/d, effect on ischaemic events	Decreased number of transient ischaemic attacks with combined treatment compared with acetylsalicylic acid alone	(73)





**Table 2.** *Continued*

Vitamin E congener	Model	Treatment and dose	Time	Experiment	Results	Reference
RRR- $\alpha$ T acetate	Rats	100 mg/100 g BW/d injected intramuscularly	7 d	Rats rendered vitamin K deficient by intraperitoneal injections of warfarin (0.01 mg/100 g BW)	Compared with warfarin alone, $\alpha$ T: Active prothrombin coagulation activity $\downarrow$ Combined measure (precursor + active protein) $\leftrightarrow$	(76)
Vitamin E (congener not specified)	Patients on warfarin therapy; <i>n</i> 25	800 or 1200 IU/d	1 month		Prothrombin coagulation activity $\leftrightarrow$	(77)
All rac- $\alpha$ T acetate	Patients on warfarin therapy; <i>n</i> 12	100 or 400 mg/d	4 weeks		Prothrombin coagulation activity was slightly reduced but non-significantly compared with pre-treatment	(76)
RRR- $\alpha$ T	Healthy males; <i>n</i> 32	667 mg/d	12 weeks		PIVKA-II $\uparrow$	(78)
$\alpha$ T (stereochemistry not specified)	Breast cancer cell lines MCF-7 and T47D	10 $\mu$ mol/l	24 h	Influence of the combination of tamoxifen and $\alpha$ T on cell proliferation	$\alpha$ T decreased the inhibitory effect of tamoxifen on cell proliferation	(86)
$\alpha$ T acetate	Breast cancer patients; <i>n</i> 7	400 mg/d	30 d	Influence of the combination of tamoxifen and $\alpha$ T on tamoxifen blood levels and markers of oestrogen stimulation in breast biopsies (pre- and post-supplementation)	Five (out of seven) patients had reduced tamoxifen concentrations; in four of these tamoxifen decreased to subtherapeutic concentrations Six (out of seven) biomarkers of oestrogen stimulation (oestrogen receptor, p-ERK) $\uparrow$ in biopsies comparing pre- and post-supplementation	(87)
RRR- $\alpha$ T	THP1 monocytes	50 $\mu$ mol/l	24 h	THP1 monocytes were treated with ritonavir alone or in combination with RRR- $\alpha$ T	Ritonavir: CD36 protein $\uparrow$ Ritonavir + $\alpha$ T: CD36 protein $\downarrow$	(95)
RRR- $\alpha$ T acetate	Rats	1.7 mg (deficient) v. 60 mg/kg diet (sufficient)	$\geq$ 9 months	Gene chip microarray expression analysis	CD36 mRNA in deficient animals v. adequate $\uparrow$	(97)
RRR- $\alpha$ T and RRR- $\gamma$ T	Rats	< 1 mg $\alpha$ T and < 1 mg $\gamma$ T (deficient) v. 12 mg $\alpha$ T and 24 mg $\gamma$ T/kg diet (sufficient)	6 months	Quantitative PCR expression analysis	CD36 mRNA in deficient animals v. adequate $\uparrow$	(98)
RRR- $\alpha$ T	Guinea-pigs	250 mg/kg diet	6 weeks	Expression of hepatic CD36 protein in response to high-fat diet and supplementation of $\alpha$ T	High fat diet v. normal: CD36 protein $\uparrow$ High-fat diet + $\alpha$ T: CD36 protein $\downarrow$	(54)

T, tocopherol;  $\downarrow$ , decrease; all rac, all racemic; PMA, phorbol 12-myristate 13-acetate;  $\leftrightarrow$ , no change; PKC, protein kinase C; BW, body weight; PIVKA-II, protein induced by vitamin K absence factor II;  $\uparrow$ , increase; p-ERK, phosphorylated extracellular signal-regulated kinase.

proteins (for example, P-glycoprotein (P-gp), also known as ATP-binding cassette transporter B1 (ABCB1) and multidrug resistance protein 1 (MDR1))<sup>(26)</sup>, may limit the intestinal absorption of drugs. Intestinal P-gp is expressed on the apical surface of enterocytes<sup>(27)</sup> and limits the bio-availability and ultimately the activity of its substrates by increasing their efflux back into the intestinal lumen<sup>(28)</sup>. A prominent example for a P-gp-mediated drug interaction is the herbal remedy St John's wort, which induces P-gp and thereby reduces the plasma concentrations of the orally administered heart medication digoxin<sup>(29)</sup>.

All four T<sub>3</sub>, but none of the T, induced P-gp mRNA expression via activation of the transcription factor pregnane X receptor (PXR; also known as steroid and xenobiotic sensing nuclear receptor) in the human colorectal adenocarcinoma cell line LS180 (incubated with 10 μmol/l of the respective T or T<sub>3</sub> for 24 h)<sup>(30)</sup>. Consistent with the effects on mRNA, P-gp protein expression and export of the P-gp substrate rhodamine 123 were increased in LS180 cells incubated with γT<sub>3</sub> (25 μmol/l; 48 h)<sup>(31)</sup>. No studies on effects of any of the remaining vitamin E congeners on intestinal P-gp protein expression or activity have been reported in the literature.

These *in vitro* studies suggest that, albeit at very high concentrations, T<sub>3</sub> may potentially enhance the transfer activity of P-gp in intestinal cells. However, the scientific literature does not contain reports from animal or human trials suggesting that intestinal P-gp-mediated interactions of vitamin E with drugs may actually occur. Thus, controlled *in vivo* trials are now warranted to elucidate the biological relevance of this potential mechanism.

**Vitamin E and pre-systemic drug metabolism.** The intestine is responsible for the uptake of nutrients, but also serves as a barrier against ingested xenobiotics<sup>(32)</sup>. Intestinal phase I enzymes, such as cytochrome P<sub>450</sub> mixed-function mono-oxygenases (CYP), and phase II enzymes, such as UDP-glucuronosyltransferases (UGT), glutathione-S-transferases (GST) or NAD(P)H:quinone oxidoreductase (NQO1), to name only a few, play a key role in the first-pass metabolism of drugs, which determines their and their metabolites' plasma concentrations. CYP3A4 is a phase I enzyme involved in the metabolism of approximately 50 % of all prescription drugs<sup>(33)</sup> and the most abundant CYP in the intestine<sup>(34)</sup>. Inhibition of CYP3A4, for example, by drinking a glass of grapefruit juice, reduces the oral availability of the sedative and CYP3A4 substrate<sup>(35)</sup> midazolam in humans<sup>(36)</sup>.

In LS180 colon cells, neither T nor T<sub>3</sub> (10 μmol/l; 24 h) altered the mRNA expression of CYP3A4<sup>(30)</sup>. In agreement with these *in vitro* findings, the intestinal protein expression of CYP3A4 did not differ between guinea-pigs fed RRR-αT at normal dietary (20 mg/kg diet) or pharmacological doses (250 mg/kg diet) for 6 weeks<sup>(37)</sup>.

During phase I metabolism, reactive quinones and epoxides may be generated. These reactive metabolites are inactivated by phase II enzymes, including NQO1,

which reduces quinones to less reactive hydroquinones, and GST, which conjugates electrophiles with glutathione. In the colon cancer cell line Colo205, αT dose-dependently (0.0001–10 μmol/l; 48 h) increased NQO1 activity, whereas GST activity remained unchanged<sup>(38)</sup>.

The phase II enzyme UGT1A conjugates lipophilic substances with UDP-glucuronic acid. UGT1A1 is the quantitatively most important UGT in the small intestine in humans and its activity in intestinal microsomes is higher than in hepatic microsomes. Hence, intestinal glucuronidation contributes significantly to the first-pass metabolism and low bioavailability of UGT1A1 substrates<sup>(39)</sup>. In LS180 cells, UGT1A1 mRNA increased at most 2-fold upon incubation with αT<sub>3</sub>, βT<sub>3</sub> or δT<sub>3</sub> (10 μmol/l; 24 h), while neither γT<sub>3</sub> nor any of the T altered UGT1A1 mRNA<sup>(30)</sup>. Such small increases in UGT1A1 mRNA, however, are unlikely to have a significant impact on UGT1A1 protein expression and/or activity. In agreement with this notion, the mean total UGT activity in the small and large intestine of rats fed 200 mg αT/kg diet for 2 weeks was unchanged relative to control animals<sup>(40)</sup>.

Thus, there is no evidence from *in vivo* studies to suggest that T or T<sub>3</sub> may alter the intestinal metabolism of drugs even though modest effects on NQO1 and UGT1A1 activity have been observed in cultured colonic adenocarcinoma cells.

### *Interactions of vitamin E with hepatic drug metabolism and excretion*

Hepatic interactions of nutrients with drugs occur predominantly on the level of the uptake of drugs into the liver and/or their conversion by hepatic phase I and II enzymes.

**Vitamin E and hepatic uptake of drugs.** Transport proteins in the plasma membrane are involved in the uptake of many drugs into hepatocytes and strongly determine their overall hepatic clearance, regardless of other subsequent reactions<sup>(41)</sup>. In humans, the organic anion transporting polypeptide (OATP C, also known as solute carrier organic anion transporter family member 1B1 (OATP1B1) or OATP2) is a liver-specific transporter facilitating the uptake of various endogenous (for example, 17β-glucuronosyl oestradiol)<sup>(42)</sup> and xenobiotic substances (for example, the drug pravastatin)<sup>(43,44)</sup> from the blood into the liver.

In rats, daily subcutaneous injections with 100 mg RRR-αT/kg body weight for 1 week, a mode of administration that was chosen to load the liver with extremely high concentrations of αT, reduced hepatic OATP C mRNA compared with control<sup>(45,46)</sup>. In humans, subcutaneous injection of αT, which has been used to treat retinopathy (retrolental fibroplasia) in prematurely born infants<sup>(47)</sup>, as well as intramuscular<sup>(48)</sup> and intravenous injections of vitamin E<sup>(49)</sup> are rare modes of administration and thus of minor relevance in the context of vitamin E–drug interactions. Also, changes in mRNA expression often do not



mirror changes in protein concentration and/or activity. In a model that more closely reflects the co-ingestion of nutrients and drugs observed in human subjects, feeding guinea-pigs for 6 weeks with either 20 or 250 mg RRR- $\alpha$ T/kg diet in the presence or absence of the OATP C substrate atorvastatin (300 mg/kg diet) did not alter hepatic OATP C protein expression, atorvastatin plasma concentrations (reflecting OATP C activity) or the lipid-lowering activity of the drug<sup>(37)</sup>.

In summary, these studies suggest that subcutaneous and potentially also intramuscular and intravenous injections with high doses of  $\alpha$ T may carry a limited risk of altering OATP C-mediated hepatic drug uptake, but there is currently no *in vivo* evidence for the biological importance of this potential mechanism of vitamin E–drug interactions. Also the oral ingestion of high doses of  $\alpha$ T does not appear to affect OATP C-mediated drug transport and efficacy *in vivo*. None of the other vitamin E congeners has been studied for impact on the hepatic uptake of drugs yet.

**Vitamin E and hepatic phase I metabolism.** During phase I metabolism in the liver, cytochrome P<sub>450</sub> enzymes (CYP) and other phase I enzymes oxidise, reduce or hydrolyse xenobiotics (such as drugs) and thereby activate them for the subsequent phase II conjugation reactions. All vitamin E congeners undergo phase I metabolism and are  $\omega$ -hydroxylated at the terminal methyl group in the side chain by a CYP, in humans probably CYP4F2<sup>(50,51)</sup>. Another member of the cytochrome family, CYP3A4, is involved in the metabolism of many drugs and can either contribute to the formation of the active agent from a parent drug or to the generation of metabolites destined for elimination<sup>(33)</sup>. Since the metabolism of all vitamin E congeners and of many drugs is catalysed by the same enzyme family, it is feasible that interactions on the level of hepatic metabolism may occur.

The impact of vitamin E on CYP3A4 expression and activity has been extensively studied in numerous *in vitro* and *in vivo* models. The expression of CYP3A4 is controlled by the nuclear receptor PXR<sup>(52)</sup>. In a reporter gene assay with HepG2 cells, T<sub>3</sub> at 10 and 50  $\mu$ mol/l activated PXR (5- and 10-fold relative to untreated control cells), while T at these concentrations were much less potent activators (maximum 2-fold induction relative to control)<sup>(30,53)</sup>. We, on the other hand, did not observe an induction of PXR at all in a HEK293 reporter gene assay with concentrations of up to 200  $\mu$ mol/l RRR- or all rac- $\alpha$ T<sup>(54)</sup>. In primary human hepatocytes, 10  $\mu$ mol/l T<sub>3</sub> induced CYP3A4 mRNA 3- to 5-fold, whereas T were ineffective<sup>(30)</sup>. Incubation of HepG2 cells with increasing concentrations (0–300  $\mu$ mol/l) of either RRR- or all rac- $\alpha$ T for 7 d dose-dependently increased mRNA expression of the orphan CYP20A1 only, but not of any of the other thirteen CYP (including CYP3A4) expressed in these cells<sup>(55)</sup>.

A number of *in vivo* studies investigated the effects of high-dose supplementation as well as vitamin E deficiency

on the expression of hepatic CYP, especially CYP3A4. In mice, feeding of a vitamin E-deficient diet (2 mg RRR- $\alpha$ T/kg diet) for 3 months reduced, whereas RRR- $\alpha$ T supplementation up to 9 months (200 mg/kg diet) did not alter hepatic mRNA expression of Cyp3a11 (the murine homologue of the human CYP3A4) compared with control (20 mg/kg diet)<sup>(56)</sup>. In a similar study, feeding rats a diet deficient or adequate in vitamin E (<2 or 60 mg RRR- $\alpha$ T) for up to 9 months did not change the mRNA of any of the thirty-three hepatic CYP expressed (including CYP3A4)<sup>(55)</sup>. On the other hand, feeding of high doses of all rac- $\alpha$ T acetate (1000 mg/kg diet) for 4 months induced the hepatic mRNA expression of Cyp3a11 approximately 1.3- to 4-fold (depending on the analytical method used) compared with control mice on a standard diet (35 mg/kg diet)<sup>(57)</sup>. However, 1000 mg/kg diet of synthetic all rac- $\alpha$ T acetate is a supraphysiological dose, which corresponds to about 7–10 g/d for an average human, and might explain the differences between the latter and the aforementioned mouse and rat studies. Although T<sub>3</sub> have a higher potential to induce CYP3A4 in reporter gene assays as well as in primary hepatocytes, hepatic mRNA expression of the murine homologue Cyp3a11 remained unchanged after oral administration of mice with  $\gamma$ T<sub>3</sub> (250  $\mu$ g/d for 7 d)<sup>(56)</sup>.

Although, as mentioned earlier, changes in mRNA may not necessarily result in similar changes in protein expression and activity, none of the above studies investigated the effects of vitamin E on CYP protein expression and activity. In an attempt to close this gap, we fed guinea-pigs with 20 or 250 mg RRR- $\alpha$ T/kg diet for 6 weeks and studied the metabolism of atorvastatin, a drug that is metabolised by CYP3A4, and its lipid-lowering efficacy. The hepatic protein expression of CYP3A4 and CYP20A1 and the serum concentrations of the CYP3A4 products of atorvastatin, *para*- and *ortho*-hydroxy-atorvastatin were similar in both groups. In agreement with this, the cholesterol-lowering activity of atorvastatin was unaltered by high-dose  $\alpha$ T feeding<sup>(37)</sup>. This is in agreement with studies in human subjects addressing the effects of  $\alpha$ T on CYP3A4-mediated drug metabolism. Patients receiving either lovastatin or simvastatin (both CYP3A4 substrates) were supplemented with 400 mg/d RRR- $\alpha$ -tocopheryl acetate for 8 weeks and CYP3A4 activity was determined by quantification of urinary cortisol and its CYP metabolite 6 $\beta$ -hydroxycortisol. In these patients, vitamin E supplementation neither altered CYP3A4 activity (urinary 6 $\beta$ -hydroxycortisol), nor the lipid-lowering activity of the statins<sup>(58)</sup>. In another human intervention trial, healthy volunteers were intravenously injected with 1 mg of the CYP3A4 substrate midazolam and its area under the plasma concentration–time curve (AUC) was determined before and after a 3-week intervention with 750 IU (503 mg) RRR- $\alpha$ T/d orally or placebo (six subjects per group). By using intravenous injection of the test compound, the investigators were able to circumvent intestinal phase I

metabolism and to investigate the effects of vitamin E on hepatic phase I metabolism only. The plasma AUC for midazolam and its metabolite hydroxymidazolam did not differ between the placebo and vitamin E groups<sup>(59)</sup>.

In summary, vitamin E–drug interactions due to altered phase I metabolism have not been reported and appear unlikely based on the available evidence. Even at high oral doses, neither T nor T<sub>3</sub> appear to significantly alter CYP activity in animals or humans.

**Vitamin E and hepatic phase II metabolism.** Phase II of xenobiotic metabolism, which takes place to a significant extent in the liver, comprises the conjugation of xenobiotics or their phase I metabolites with polar groups, such as glucuronic acid, sulfate, glutathione or amino acids, to enable rapid biliary and urinary elimination. The inhibition of phase II enzymes, such as the UGT and GST, results in increased blood and tissue concentrations of their substrates. The induction of phase II enzymes, on the other hand, may increase the clearance and thereby reduce the activity of drugs.

None of the eight vitamin E congeners (incubation with 10 μmol/l for 24 h) altered UGT1A1 mRNA in primary human hepatocytes<sup>(30)</sup> and feeding rats 200 mg αT/kg diet for 2 weeks had no effect on hepatic UGT activity<sup>(40)</sup>. A small increase (1.76– to 2.68-fold, depending on assay) in hepatic GST activity was observed in rats fed very high doses of αT (2500 mg/kg diet) for 10 d<sup>(60)</sup>. This dose, however, would be equal to more than 17 g/d of αT for a 70 kg human. Similarly, feeding mice with 1000 mg/kg diet of all rac-αT acetate for 4 months led to a small increase (maximum 1.9-fold) in the hepatic mRNA expression of the murine Gstm3 compared with mice on a standard diet (35 mg all rac-αT acetate/kg diet)<sup>(57)</sup>. In disagreement with the above-described effects *in vivo*, studies with isolated human GST P1-1 suggest that the concentrations at which T and T<sub>3</sub> inhibit GST activity by 50 % (IC<sub>50</sub>) are: αT, 0.7; δT, 0.8<sup>(61)</sup>; αT<sub>3</sub>, 1.8; and γT<sub>3</sub>, 0.7 μmol/l<sup>(62)</sup>. However, direct interactions of vitamin E with enzymes in artificial *in vitro* systems may differ significantly from cellular systems and the situation *in vivo*, where vitamin E resides in membranes and may thus not have direct access to these soluble cytosolic proteins.

The mRNA expression of another phase II enzyme, sulfotransferase 2A, was significantly induced (11-fold) in the aforementioned mouse study, where animals were fed 1000 compared with 35 mg/kg diet all rac-αT acetate for 4 months<sup>(57)</sup>.

Overall, there is no evidence to support the notion that T or T<sub>3</sub> ingested at normal dietary or supplemental doses may alter the activity of hepatic phase II metabolism in the living organism.

### Vitamin E and hepatic drug export

Some researchers consider the export of xenobiotics or their phase I and II metabolites from cells, facilitated by

the activity of membrane transporters such as the multi-drug resistance proteins (for example, P-gp) and other ATP-binding cassette transporters, as phase III of xenobiotic metabolism. In the liver, hepatocytes lining the biliary canalicular surface express P-gp<sup>(27)</sup> and secrete substrates, including vitamin E and its metabolites<sup>(63)</sup>, into the bile.

In primary human hepatocytes, P-gp mRNA expression was not affected by incubation with any of the T or T<sub>3</sub> (10 μmol/l; 24 h)<sup>(30)</sup>. In mice fed very high doses of all rac-αT acetate (1000 mg/kg diet compared with 35 mg/kg diet) for 4 months, P-gp mRNA expression increased 1.9-fold in the liver<sup>(57)</sup>. The same authors, in another *in vivo* experiment, investigated the effects of daily subcutaneous injections of two different formulations (micellised vitamin E (Emcelle) and vitamin E dissolved in 20 % ethanol and 1 % benzyl alcohol (VitalE)) of high-dose RRR-αT (100 mg/kg body weight for up to 9 d) in rats. Both formulations increased P-gp mRNA (VitalE, 3.57-fold; Emcelle, about 10-fold), while P-gp protein expression was only reported for VitalE injections, which significantly increased it 3-fold<sup>(45)</sup>.

Thus, two animal studies report that αT may induce hepatic P-gp expression when injected intraperitoneally or subcutaneously in high doses, which are unusual routes of administration and as such of limited relevance in a nutritional context. Overall, there are no publications to suggest that the consumption of T or T<sub>3</sub> at conventional dietary or supplemental doses may induce the expression of hepatic efflux transporters or cause hepatic membrane-transporter-mediated drug interactions *in vivo*. However, it cannot presently be excluded that vitamin E taken at extremely high doses may affect the expression and perhaps the activity of hepatic drug exporters.

### Vitamin E and pharmacodynamics

The effects of vitamin E on pharmacodynamics can be mediated indirectly by altering drug pharmacokinetics, as described in the previous sections, or more directly by targeting the molecular mechanisms underlying drug activity. The latter may be achieved by direct interactions with the drug's molecular targets or by actions on additional (independent) targets, such as signalling molecules, which alter the activity of a drug.

**Vitamin E and blood coagulation.** The best-known vitamin E–drug interactions are those with drugs that affect platelet aggregation and blood coagulation, which are presently the only interactions listed by the US Food and Drug Administration (FDA) in their consumer health information<sup>(64)</sup>. Blood coagulation is the process during which blood is (locally) converted from a liquid to a gel and is the body's answer to injury ensuring that ruptured blood vessels are rapidly sealed off to minimise blood loss.


In brief, coagulation is initiated when collagen, which is normally buried underneath the epithelium, is exposed

and platelets adhere to it via collagen-specific surface receptors (glycoprotein Ia/IIa). Platelet adhesion and the release of signalling molecules (including tissue factor and von Willebrand factor) from both the injured endothelium and the platelets activate further platelets and the intrinsic and extrinsic coagulation pathways. These coagulation cascades include a number of enzymes that are synthesised as precursors in the liver, where they undergo vitamin K-dependent carboxylation by  $\gamma$ -glutamyl-carboxylase, and are secreted as mature (carboxylated) proteins into the bloodstream (Fig. 3). The adhesion and aggregation of platelets at the site of injury result in the formation of a patch that is stabilised by the deposition and maturation of the fibrous protein fibrin (also known as factor Ia). Fibrin is formed by polymerisation of its precursor fibrinogen through the activity of the enzyme thrombin, which itself is activated by cleavage of prothrombin (factor II). Activated platelets secrete granules containing ADP, serotonin, platelet-activating factor, von Willebrand factor, platelet factor 4 and thromboxane A<sub>2</sub> (TXA<sub>2</sub>) that induce platelet aggregation through cross-linkage of activated glycoprotein IIb-IIIa and fibrinogen. TXA<sub>2</sub> is synthesised from arachidonic acid through the cyclo-oxygenase (COX) and hydroperoxidase catalytic activities of PG synthase, via the intermediate PGH<sub>2</sub>. The COX reaction is the rate-limiting step in the synthesis of prostaglandins and thromboxanes and is inhibited by drugs, such as acetylsalicylic acid, which thereby reduce platelet aggregation.

Platelet activation is a complex regulatory pathway with multiple agonists, including ADP, TXA<sub>2</sub>, collagen, adrenaline and many more. *Ex vivo* platelet aggregation can be activated by either endogenous agonist or phorbol

myristate acetate (PMA), a protein kinase C (PKC) activator. PKC is a family of enzymes catalysing the phosphorylation and thus controlling the activity of many regulatory proteins. PKC activation stimulates, among other things, platelet granule secretion, TXA<sub>2</sub> synthesis and thereby the aggregation of platelets<sup>(65)</sup>.

**Vitamin E and platelet adhesion, activation and aggregation.** Platelets obtained from healthy volunteers and pre-incubated with RRR- $\alpha$ T for 5 min (0.016–16.22  $\mu$ mol/l) exhibited a dose-dependent reduction in the metabolism of arachidonic acid to TXA<sub>2</sub> and PGD<sub>2</sub>. The authors suggested that this may be due to an inhibition of COX activity in  $\alpha$ T-treated platelets<sup>(66)</sup>. Incubation of platelets from healthy human subjects with supraphysiological concentrations (500  $\mu$ mol/l) of RRR- $\alpha$ T or RRR- $\alpha$ -tocopheryl acetate, but not all rac- $\alpha$ T, inhibited arachidonic acid- or PMA-induced platelet aggregation and PKC-mediated protein phosphorylation<sup>(67,68)</sup>. Although 500  $\mu$ mol/l  $\alpha$ T is a supranutritional dose, the authors observed that the cellular content of  $\alpha$ T in the incubated platelets was comparable with that of platelets isolated from human subjects receiving 800 mg  $\alpha$ T/d for 14 d<sup>(67)</sup>. In the latter study, platelets isolated from human subjects supplemented for 14 d with 267, 533 or 800 mg  $\alpha$ T/d, PMA-induced platelet aggregation was dose-dependently reduced, arachidonic acid-induced aggregation was reduced in the 800 mg/d group only, and ADP-induced aggregation was not altered by  $\alpha$ T<sup>(67)</sup>. PKC-mediated phosphorylation was determined in two subjects and was inhibited after 2 weeks' supplementation with 800 mg RRR  $\alpha$ T/d compared with baseline<sup>(67)</sup>. In another study where healthy volunteers were given placebo, 100 mg all rac- $\alpha$ T acetate, or mixed T (20 mg RRR- $\alpha$ T, 100 mg



**Fig. 3.** (a) The precursors of mature factor II, VII, IX and X are carboxylated by  $\gamma$ -glutamyl-carboxylase to their mature forms. The cofactor (vitamin K) for this reaction is oxidised and then regenerated by vitamin K epoxidase. The inhibition of vitamin K epoxidase by, for example, warfarin decreases the recycling of oxidised vitamin K and results in a shortage of the cofactor for the carboxylation reaction. (b) Upon endothelial injury, platelets bind to sub-endothelial collagen (via glycoprotein Ia/IIa (GPIIa/IIa) and von Willebrand factor (vWF)). Tissue factor (TF) is also released to activate the extrinsic pathway. Interaction between GPVI and collagen induces cyclo-oxygenase (COX) and triggers the formation of thromboxane A<sub>2</sub> (TXA<sub>2</sub>), which cross-links platelets via GPIIb-IIIa and fibrinogen. Protein kinase C (PKC) activates, among other things, TXA<sub>2</sub> formation and granule secretion. (A colour version of this figure can be found online at [www.journals.cambridge.org/nrr](http://www.journals.cambridge.org/nrr)).

RRR- $\gamma$ T and 40 mg RRR- $\delta$ T) daily for 8 weeks, only mixed T significantly reduced ADP-induced, but not PMA-induced platelet aggregation, although phosphorylation of PKC was significantly inhibited by both vitamin E interventions<sup>(69)</sup>. In yet another study, platelet adhesion to collagen was dose-dependently reduced in twelve healthy subjects supplemented with increasing doses of all rac- $\alpha$ T acetate (400, 800 and 1200 mg/d for 2 weeks, each; 6 weeks supplementation in total) compared with eleven healthy control subjects. Platelet aggregation (induced with collagen, adrenaline or ADP) was only modestly decreased in these subjects and only collagen-induced platelet aggregation was significantly reduced ( $P < 0.05$ ) in women, but not men<sup>(70)</sup>. The same study also analysed the combination of all rac- $\alpha$ T with the analgesic, antipyretic and anti-inflammatory drug acetylsalicylic acid (aspirin)<sup>(70)</sup>, which itself inhibits the COX-catalysed formation of TXA<sub>2</sub> from arachidonic acid<sup>(71)</sup>. The combined oral administration of all rac- $\alpha$ T acetate in doses up to 1200 mg/d with 300 mg aspirin every other day had no effect on collagen-, adrenaline- or ADP-induced platelet aggregation or platelet adhesion to collagen compared with vitamin E or acetylsalicylic acid alone<sup>(70)</sup>. In a subset of the Alpha-Tocopherol Beta-Carotene Cancer Prevention Study, male smokers aged 50–59 years who received a combination of 100–3200 mg acetylsalicylic acid and 50 mg all rac- $\alpha$ T acetate/d and were followed for 5–7 years showed a significant increase in oral 'bleeding on probing' (gingival bleeding) compared with subjects receiving acetylsalicylic acid alone<sup>(72)</sup>. Accordingly, in patients who previously suffered from an ischaemic event, the combination of acetylsalicylic acid (325 mg) and all rac- $\alpha$ T acetate (400 mg/d) for up to 2 years ( $n$  52), compared with acetylsalicylic acid alone ( $n$  48), was more effective in the prevention of transient ischemic attacks, suggesting a synergistic activity of  $\alpha$ T with aspirin<sup>(73)</sup>.

The mechanism for the reduction in *ex vivo* platelet aggregation observed in the above studies still remains to be fully elucidated. The inhibition of PKC by  $\alpha$ T may be an important factor, since platelet aggregation induced by the PKC activator PMA was reduced by vitamin E in most studies concomitantly with a reduction in PKC activity<sup>(67–69)</sup>. Interactions with vitamin K-dependent coagulation cascades, however, may be part of the explanation, as discussed in the next section.

**Vitamin E, vitamin K and the extrinsic and intrinsic coagulation pathways.** As mentioned earlier, the coagulation factors II, VII, IX and X undergo post-translational carboxylation by  $\gamma$ -glutamyl-carboxylase (Fig. 3), an enzyme that requires vitamin K as a cofactor, in the liver. Vitamin K is oxidised in the carboxylation reaction and requires subsequent reduction by vitamin K epoxide reductase in order to take part in another cycle of carboxylation<sup>(74)</sup>. Some anticoagulants, such as warfarin, act by reducing the vitamin K epoxide reductase-catalysed recycling of oxidised vitamin K, thereby limiting the

carboxylation of the precursors of factors II (prothrombin), VII, IX and X to their mature forms (Fig. 3)<sup>(75)</sup>.

In rats with decreased concentrations of reduced vitamin K (induced by intraperitoneal injections of warfarin), intramuscular injections of RRR- $\alpha$ T acetate (1000 mg/kg body weight/d for 7 d) reduced mature prothrombin coagulant activity compared with animals treated with warfarin only, but not the amount of the prothrombin precursor produced in the liver<sup>(76)</sup>. Since  $\alpha$ T influences the formation of the mature prothrombin rather than the production of the precursor, this might be caused by a vitamin E-induced aggravation of vitamin K deficiency. Prothrombin activity in warfarin-treated patients supplemented with vitamin E (800 or 1200 IU vitamin E (congener not specified)/d,  $n$  13; or 100 or 400 IU all rac  $\alpha$ T-acetate/d,  $n$  6) for 4 weeks did not differ significantly from that of placebo-treated patients<sup>(76,77)</sup>. Although these trials are limited due to their small sample size, they suggest that doses up to 1200 IU vitamin E have no clinical effects in patients on warfarin therapy. Nevertheless, vitamin E may interfere with vitamin K metabolism to some extent. Human subjects supplemented with 1000 IU RRR- $\alpha$ T/d (12 weeks) showed an increase in PIVKA-II (protein induced by vitamin K absence factor II) plasma concentrations<sup>(78)</sup>, which is a marker of subclinical vitamin K deficiency<sup>(79)</sup>. Interactions between the metabolism of vitamins E and K are feasible, since both appear to share the same metabolic fate and may be  $\omega$ -hydroxylated by the same enzyme(s) (*inter alia* CYP4F2)<sup>(50,51,80)</sup>. However, vitamin K  $\omega$ -hydroxylation was neither affected in cells incubated with, nor in animals fed high doses of, vitamin E<sup>(46,81)</sup>. Thus, the mechanism of vitamin E-induced changes in vitamin K status remains unclear and warrants further investigation.

In summary, reduced blood coagulation and increased bleeding have been observed in some cases in animals and human subjects treated with aspirin or warfarin simultaneously with high doses of vitamin E (mainly  $\alpha$ T). Such a synergistic activity may be desirable, for example in patients at risk for CVD or stroke, or undesirable, as in the case of increased (gastrointestinal) bleeding, a known side effect of aspirin<sup>(82)</sup>. Vitamin K status and coagulation activity should be monitored in patients supplementing vitamin E while on warfarin therapy (or taking other medication interfering with vitamin K-dependent activation of coagulation factors) to detect and counteract changes in haemostasis.

#### *Interactions of vitamin E of an unknown kind that alter drug efficacy*

**Vitamin E may impair tamoxifen drug activity in breast cancer patients.** Oestrogen-responsive cancer may be treated with the selective oestrogen receptor modulator tamoxifen<sup>(83)</sup>, but most patients eventually develop resistance to the drug<sup>(84)</sup>. Tamoxifen rapidly increases intracellular concentrations of the second messenger Ca<sup>2+</sup><sup>(85)</sup>.



Cell-culture studies suggest that vitamin E may alter the activity of tamoxifen. In oestrogen-responsive mammary carcinoma cells (MCF-7, T47D), incubation with tamoxifen in the presence of  $\alpha$ T (10  $\mu$ mol/l; 24 h) inhibited the tamoxifen-induced rise in intracellular  $\text{Ca}^{2+}$  and reduction in cell proliferation<sup>(86)</sup>. In a small prospective study with seven breast cancer patients on tamoxifen therapy, supplementation with 400 mg/d  $\alpha$ T acetate for 30 d reduced tamoxifen blood concentrations in five patients and increased markers of tamoxifen resistance in six of the seven patients. In four of these patients, tamoxifen blood concentrations dropped to below the minimum therapeutic concentration<sup>(87)</sup>.

In summary, the limited evidence from cell-culture and human intervention trials suggests that it might be undesirable for women on tamoxifen therapy to consume dietary supplements containing high doses of  $\alpha$ T, as they may impair drug efficacy. However, before valid conclusions can be drawn, sufficiently powered randomised clinical trials investigating the interactions of  $\alpha$ T with tamoxifen pharmacokinetics and pharmacodynamics are required.

*High-dose vitamin E supplementation may alter the pharmacokinetics of cyclosporine A.* Cyclosporine A is an immunosuppressant administered to prevent graft rejection in organ transplant recipients. One of the side effects of cyclosporine A is an increased generation of reactive oxygen species, which was suggested to be involved in cyclosporine A-induced nephrotoxicity<sup>(88)</sup>. To investigate if supplementation with antioxidants may be beneficial in cyclosporine A-treated renal transplant recipients, ten patients were supplemented with an antioxidant cocktail (800 IU vitamin E ( $\alpha$ T), 1000 mg vitamin C and 6 mg  $\beta$ -carotene per d) or placebo for 6 months in a cross-over study with a 6-month washout period<sup>(89)</sup>. All cyclosporine A trials reported in this paragraph used  $\alpha$ T in the dietary supplements (ML Blackhall, RG Fassett, JE Sharman, DP Geraghty and JS Coombes, personal communication). The antioxidant cocktail did not alter any of the markers of oxidative stress, but decreased cyclosporine A concentrations in blood by 24 %<sup>(89)</sup>. In a comparable trial, kidney transplant recipients on cyclosporine A therapy were supplemented for 3 months with placebo ( $n$  28) or 1000 mg vitamin C plus 300 mg vitamin E ( $\alpha$ T;  $n$  25) per d and reduced cyclosporine A blood concentrations were reported for the vitamin-supplemented group<sup>(90)</sup>. Similar observations were made retrospectively in heart transplant patients; compared with baseline, cyclosporine A blood concentrations decreased by 30 % when patients were supplemented with 1000 mg vitamin C and 800 IU vitamin E ( $\alpha$ T) per d<sup>(91)</sup>. However, in all studies, vitamin E was administered in combination with vitamin C, which makes it impossible to discern the contribution of each individual vitamin.

The effect of vitamin E alone on cyclosporine A pharmacokinetics was studied earlier in healthy volunteers before and after supplementation with 800 IU vitamin E ( $\alpha$ T) per d

for 6 weeks. Vitamin E supplementation had no impact on the glomerular filtration rate or peak plasma concentration, but reduced the AUC of a single oral dose of cyclosporine A (5 mg) by 21 % compared with baseline<sup>(92)</sup>.

The mechanism underlying the interaction of vitamin E with cyclosporine A is currently unknown. In the trials discussed above, vitamin E and cyclosporine A were administered separately, which eliminates the possibility of a direct interaction. Cyclosporine A is a substrate for intestinal P-gp and CYP3A4 as well as hepatic CYP3A4 and CYP3A5<sup>(93)</sup>, which makes pre-systemic interactions on the level of drug absorption and intestinal and systemic interactions possible. Induction of intestinal and hepatic metabolism would both decrease the plasma concentration and AUC of cyclosporine A. However, as discussed above, orally administered  $\alpha$ T does not induce the expression of CYP or membrane transporters<sup>(37,55,56,58)</sup>, rendering this mode of interaction rather unlikely.

Since vitamin E supplementation at daily doses of  $\geq$  300 mg affected the pharmacokinetics of cyclosporine A in four independent human trials, patients treated with this drug should abstain from the consumption of dietary supplements containing such high doses of vitamin E or at least have the blood concentration and efficacy of the drug tested regularly by their physician.

*$\alpha$ -Tocopherol supplementation reverses the over-expression of the fatty acid translocase CD36.* The potential interaction of vitamin E with ritonavir described below differs from the cases discussed above in that it relates to a potential adverse interaction in the case of vitamin E deficiency rather than under vitamin E supplementation. Ritonavir is a drug that specifically inhibits the HIV 1 protease and thereby HIV replication<sup>(94)</sup> and is used for the treatment of HIV-infected patients. Adverse effects observed upon long-term treatment with ritonavir include hyperlipidaemia and accelerated atherosclerosis, which may in part be caused by an over-expression of the scavenger receptor CD36 protein in macrophages<sup>(95)</sup>.

In THP1 monocytes, ritonavir-induced CD36 (over-) expression was reversed by incubation with RRR- $\alpha$ T (50  $\mu$ mol/l; 24 h)<sup>(96)</sup>. Further evidence to support a role of vitamin E in the expression of CD36 came from two studies in rats fed vitamin E-deficient (1.7 mg  $\alpha$ T acetate; <1 mg total T/kg diet, respectively) compared with vitamin E-sufficient (60 mg  $\alpha$ T acetate; 12 mg RRR- $\alpha$ T and 24 mg RRR- $\gamma$ T/kg, respectively) diets for 6 and up to 9 months, respectively, in which an up-regulation of hepatic CD36 mRNA was observed in the deficient animals from the third month onwards<sup>(97,98)</sup>. In guinea-pigs fed a high-fat diet (21 % fat by weight), CD36 protein expression in the liver was almost twice that observed in control animals on a 5 % fat diet. Inclusion of high doses of RRR- $\alpha$ T (250 mg/kg diet) in the high-fat diet reversed the increase in hepatic CD36 expression back to the expression level observed in animals fed the 5 % fat control diet<sup>(54)</sup>.

Taken together, these data suggest that vitamin E deficiency might increase CD36 expression and could thus potentially aggravate ritonavir-induced hyperlipidaemia. Therefore, a sufficient intake (and perhaps even low-dose supplementation) of vitamin E may be beneficial in ritonavir-treated patients. However, there are presently no *in vivo* data available to corroborate that such an interaction may be biologically relevant in humans and further studies addressing the potential benefit as well as the safety of a co-administration of ritonavir and vitamin E are necessary before valid recommendations can be given.

## Conclusions

There is no convincing evidence from animal models or controlled human intervention trials to suggest that T–drug interactions may be caused by an induction of xenobiotic metabolism or trans-membrane transport. Corresponding *in vivo* studies for T<sub>3</sub> are presently not available and are therefore warranted. Based on the documented safety of αT in randomised clinical trials<sup>(99,100)</sup>, the FDA has set its tolerable upper intake level for adults at 1000 mg/d<sup>(1)</sup>. There is little evidence to suggest potent health benefits from the intake of such high doses in healthy individuals; specific subgroups of the population (for example, type 2 diabetics with a hapto-globin 2–2 genotype), however, may benefit from αT supplementation at doses below 300 mg/d<sup>(21,101–103)</sup>. The dietary intake of T and T<sub>3</sub> and the consumption of medium-dose supplements (<300 mg/d) appear safe and unlikely to cause adverse drug interactions and are in accordance with the tolerable upper intake levels for adults set by the European Food Safety Authority (300 mg/d)<sup>(104)</sup>. Consumption of high-dose vitamin E supplements may lead to interactions with the drugs aspirin, warfarin, tamoxifen and cyclosporine A and may alter their activities. Patients treated with these drugs should therefore consult their physician and have drug efficacy tested if they wish to consume vitamin E supplements. For the majority of drugs, however, interactions with vitamin E, even at high doses, are unlikely and have not been observed.

## Acknowledgements

There are no conflicts of interest.

## References

- Hathcock JN, Azzi A, Blumberg J, *et al.* (2005) Vitamins E and C are safe across a broad range of intakes. *Am J Clin Nutr* **81**, 736–745.
- Miller ER III, Pastor-Barriuso R, Dalal D, *et al.* (2005) Meta-analysis: high-dosage vitamin E supplementation may increase all-cause mortality. *Ann Intern Med* **142**, 37–46.
- Bjelakovic G, Nikolova D, Gluud LL, *et al.* (2007) Mortality in randomized trials of antioxidant supplements for primary and secondary prevention: systematic review and meta-analysis. *JAMA* **297**, 842–857.
- Brigelius-Flohe R (2003) Vitamin E and drug metabolism. *Biochem Biophys Res Commun* **305**, 737–740.
- Frank J, Chin XWD, Schrader C, *et al.* (2012) Do tocotrienols have potential as neuroprotective dietary factors? *Ageing Res Rev* **11**, 163–180.
- Kamal-Eldin A & Appelqvist LA (1996) The chemistry and antioxidant properties of tocopherols and tocotrienols. *Lipids* **31**, 671–701.
- Evans HM & Bishop KS (1922) On the existence of a hitherto unrecognized dietary factor essential for reproduction. *Science* **56**, 650–651.
- Leth T & Sondergaard H (1977) Biological activity of vitamin E compounds and natural materials by the resorption-gestation test, and chemical determination of the vitamin E activity in foods and feeds. *J Nutr* **107**, 2236–2243.
- Burton GW, Joyce A & Ingold KU (1982) First proof that vitamin E is major lipid-soluble, chain-breaking antioxidant in human blood plasma. *Lancet* **ii**, 327.
- Azzi A, Gysin R, Kempna P, *et al.* (2004) Vitamin E mediates cell signaling and regulation of gene expression. *Ann N Y Acad Sci* **1031**, 86–95.
- Rimbach G, Minihane AM, Majewicz J, *et al.* (2002) Regulation of cell signalling by vitamin E. *Proc Nutr Soc* **61**, 415–425.
- Frank J, de Pascual Teresa S & Rimbach G (2006) Nutrigenomics – new frontiers in antioxidant research. *Food Sci Technol Bull: Functional Foods* **3**, 1–12.
- Grebenstein N, Schumacher M, Graeve L, *et al.* (2014) α-Tocopherol transfer protein is not required for the discrimination against γ-tocopherol *in vivo* but protects it from side-chain degradation *in vitro*. *Mol Nutr Food Res* **58**, 1052–1060.
- Institute of Medicine (2000) Vitamin E. In *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*, pp. 186–283. Washington, DC: The National Academies Press.
- Bundesministerium für Ernährung, Landwirtschaft und Verbraucherschutz (Federal Ministry of Food, Agriculture and Consumer Protection) (2008) Ergebnisbericht, Teil 2; Nationale Verzehrsstudie II (Results Report, Part 2; National Nutrition Survey II) [http://www.mri.bund.de/fileadmin/Institute/EV/NVSII\\_Abschlussbericht\\_Teil\\_2.pdf](http://www.mri.bund.de/fileadmin/Institute/EV/NVSII_Abschlussbericht_Teil_2.pdf) (accessed July 2014).
- Schwab S, Heier M, Schneider A, *et al.* (2014) The use of dietary supplements among older persons in Southern Germany – results from the KORA-age study. *J Nutr Health Aging* **18**, 510–519.
- Balluz LS, Kieszak SM, Philen RM, *et al.* (2000) Vitamin and mineral supplement use in the United States. Results from the third National Health and Nutrition Examination Survey. *Arch Fam Med* **9**, 258–262.
- Wolfram G (2003) New reference values for nutrient intake in Germany, Austria and Switzerland (DACH-Reference Values). *Forum Nutr* **56**, 95–97.
- Deutsche Gesellschaft für Ernährung e.V. (DGE), Österreichische Gesellschaft für Ernährung (ÖGE) & Schweizerische Gesellschaft für Ernährungsforschung (SGE) (2013) *Referenzwerte Für Die Nährstoffzufuhr*. 1. Auflage ed. (*Dietary Reference Values*. 1st edition). Neustadt an der Weinstraße: Umschau Buchverlag.
- Bell SJ & Grochoski GT (2008) How safe is vitamin E supplementation? *Crit Rev Food Sci Nutr* **48**, 760–774.
- Frank J & Rimbach G (2009) Vitamin E in disease prevention – a critical appraisal of vitamin E supplementation trials. *Aktuel Ernaehr Med* **34**, 131–140.





22. Brigelius-Flohe R (2007) Adverse effects of vitamin E by induction of drug metabolism. *Genes Nutr* **2**, 249–256.
23. Benet L (1984) Pharmacokinetics: basic principles and its use as a tool in drug metabolism. In *Drug Metabolism and Drug Toxicity*, pp. 199–211 [R Mitchell and M Horning, editors]. New York: Raven Press.
24. Chan LN (2002) Drug–nutrient interaction in clinical nutrition. *Curr Opin Clin Nutr Metab Care* **5**, 327–332.
25. Buettner GR (1993) The pecking order of free radicals and antioxidants: lipid peroxidation,  $\alpha$ -tocopherol, and ascorbate. *Arch Biochem Biophys* **300**, 535–543.
26. Roninson IB, Chin JE, Choi KG, *et al.* (1986) Isolation of human MDR DNA sequences amplified in multidrug-resistant KB carcinoma cells. *Proc Natl Acad Sci U S A* **83**, 4538–4542.
27. Thiebaut F, Tsuruo T, Hamada H, *et al.* (1987) Cellular localization of the multidrug-resistance gene product P-glycoprotein in normal human tissues. *Proc Natl Acad Sci U S A* **84**, 7735–7738.
28. Mayer U, Wagenaar E, Beijnen JH, *et al.* (1996) Substantial excretion of digoxin via the intestinal mucosa and prevention of long-term digoxin accumulation in the brain by the *mdr 1a* P-glycoprotein. *Br J Pharmacol* **119**, 1038–1044.
29. Dürr D, Stieger B, Kullak-Ublick GA, *et al.* (2000) St John's wort induces intestinal P-glycoprotein/MDR1 and intestinal and hepatic CYP3A4. *Clin Pharmacol Ther* **68**, 598–604.
30. Zhou C, Tabb MM, Sadatrafiei A, *et al.* (2004) Tocotrienols activate the steroid and xenobiotic receptor, SXR, and selectively regulate expression of its target genes. *Drug Metab Dispos* **32**, 1075–1082.
31. Abuznait AH, Qosa H, O'Connell ND, *et al.* (2011) Induction of expression and functional activity of P-glycoprotein efflux transporter by bioactive plant natural products. *Food Chem Toxicol* **49**, 2765–2772.
32. Kaminsky LS & Zhang Q (2003) The small intestine as a xenobiotic-metabolizing organ. *Drug Metab Dispos* **31**, 1520–1525.
33. Guengerich FP (1999) Cytochrome P-450 3A4: regulation and role in drug metabolism. *Annu Rev Pharmacol Toxicol* **39**, 1–17.
34. Benet LZ & Cummins CL (2001) The drug efflux-metabolism alliance: biochemical aspects. *Adv Drug Deliv Rev* **50**, S3–S11.
35. Paine MF, Shen DD, Kunze KL, *et al.* (1996) First-pass metabolism of midazolam by the human intestine. *Clin Pharmacol Ther* **60**, 14–24.
36. Kupferschmidt HH, Ha HR, Ziegler WH, *et al.* (1995) Interaction between grapefruit juice and midazolam in humans. *Clin Pharmacol Ther* **58**, 20–28.
37. Podszun MC, Grebenstein N, Hofmann U, *et al.* (2013) High-dose supplementation with natural  $\alpha$ -tocopherol does neither alter the pharmacodynamics of atorvastatin nor its phase I metabolism in guinea pigs. *Toxicol Appl Pharmacol* **266**, 452–458.
38. Wang W & Higuchi CM (1995) Induction of NAD(P)H:quinone reductase by vitamins A, E and C in Colo205 colon cancer cells. *Cancer Lett* **98**, 63–69.
39. Fisher MB, Vandenbranden M, Findlay K, *et al.* (2000) Tissue distribution and interindividual variation in human UDP-glucuronosyltransferase activity: relationship between UGT1A1 promoter genotype and variability in a liver bank. *Pharmacogenetics* **10**, 727–739.
40. Van der Logt EM, Roelofs HM, van Lieshout EM, *et al.* (2004) Effects of dietary anticarcinogens and nonsteroidal anti-inflammatory drugs on rat gastrointestinal UDP-glucuronosyltransferases. *Anticancer Res* **24**, 843–850.
41. Shitara Y, Sato H & Sugiyama Y (2005) Evaluation of drug–drug interaction in the hepatobiliary and renal transport of drugs. *Annu Rev Pharmacol Toxicol* **45**, 689–723.
42. König J, Cui Y, Nies AT, *et al.* (2000) A novel human organic anion transporting polypeptide localized to the basolateral hepatocyte membrane. *Am J Physiol Gastrointest Liver Physiol* **278**, G156–G164.
43. Hsiang B, Zhu Y, Wang Z, *et al.* (1999) A novel human hepatic organic anion transporting polypeptide (OATP2) identification of a liver-specific human organic anion transporting polypeptide and identification of rat and human hydroxymethylglutaryl-CoA reductase inhibitor transporters. *J Biol Chem* **274**, 37161–37168.
44. Hagenbuch B & Meier PJ (2003) The superfamily of organic anion transporting polypeptides. *Biochim Biophys Acta* **1609**, 1–18.
45. Traber MG, Labut EM, Leonard SW, *et al.* (2011)  $\alpha$ -Tocopherol injections in rats up-regulate hepatic ABC transporters, but not cytochrome P<sub>450</sub> enzymes. *Free Radic Biol Med* **51**, 2031–2040.
46. Farley SM, Leonard SW, Labut EM, *et al.* (2012) Vitamin E decreases extra-hepatic menaquinone-4 concentrations in rats fed menadione or phyloquinone. *Mol Nutr Food Res* **56**, 912–922.
47. Bougle D, Boutroy MJ, Heng J, *et al.* (1986) Plasma kinetics of parenteral tocopherol in premature infants. *Dev Pharmacol Ther* **9**, 310–316.
48. Guggenheim MA, Ringel SP, Silverman A, *et al.* (1982) Progressive neuromuscular disease in children with chronic cholestasis and vitamin E deficiency: diagnosis and treatment with  $\alpha$ -tocopherol. *J Pediatr* **100**, 51–58.
49. Brion LP, Bell EF, Raghuvver TS, *et al.* (2004) What is the appropriate intravenous dose of vitamin E for very-low-birth-weight infants? *J Perinatol* **24**, 205–207.
50. Sontag TJ & Parker RS (2002) Cytochrome P<sub>450</sub>  $\omega$ -hydroxylase pathway of tocopherol catabolism. Novel mechanism of regulation of vitamin E status. *J Biol Chem* **277**, 25290–25296.
51. Bardowell SA, Duan F, Manor D, *et al.* (2012) Disruption of mouse cytochrome P<sub>450</sub> 4f14 (Cyp4f14) causes severe perturbations in vitamin E metabolism. *J Biol Chem* **287**, 26077–26086.
52. Lehmann JM, McKee DD, Watson MA, *et al.* (1998) The human orphan nuclear receptor PXR is activated by compounds that regulate CYP3A4 gene expression and cause drug interactions. *J Clin Invest* **102**, 1016–1023.
53. Landes N, Pfluger P, Kluth D, *et al.* (2003) Vitamin E activates gene expression via the pregnane X receptor. *Biochem Pharmacol* **65**, 269–273.
54. Podszun MC, Grebenstein N, Spruss A, *et al.* (2014) Dietary  $\alpha$ -tocopherol and atorvastatin reduce high-fat-induced lipid accumulation and down-regulate CD36 protein in the liver of guinea pigs. *J Nutr Biochem* **25**, 573–579.
55. Hundhausen C, Frank J, Rimbach G, *et al.* (2006) Effect of vitamin E on cytochrome P<sub>450</sub> mRNA levels in cultured hepatocytes (HepG2) and in rat liver. *Cancer Genom Proteom* **3**, 183–190.
56. Kluth D, Landes N, Pfluger P, *et al.* (2005) Modulation of Cyp3a11 mRNA expression by  $\alpha$ -tocopherol but not  $\gamma$ -tocotrienol in mice. *Free Radic Biol Med* **38**, 507–514.
57. Mustacich DJ, Gohil K, Bruno RS, *et al.* (2009)  $\alpha$ -Tocopherol modulates genes involved in hepatic xenobiotic pathways in mice. *J Nutr Biochem* **20**, 469–476.
58. Leonard SW, Joss JD, Mustacich DJ, *et al.* (2007) Effects of vitamin E on cholesterol levels of hypercholesterolemic patients receiving statins. *Am J Health Syst Pharm* **64**, 2257–2266.



59. Clarke MW, Burnett JR, Wu JH, *et al.* (2009) Vitamin E supplementation and hepatic drug metabolism in humans. *J Cardiovasc Pharmacol* **54**, 491–496.
60. Manson MM, Ball H, Barrett MC, *et al.* (1997) Mechanism of action of dietary chemoprotective agents in rat liver: induction of phase I and II drug metabolizing enzymes and aflatoxin B1 metabolism. *Carcinogenesis* **18**, 1729–1738.
61. Van Haaften RI, Evelo CT, Penders J, *et al.* (2001) Inhibition of human glutathione-S-transferase P1-1 by tocopherols and  $\alpha$ -tocopherol derivatives. *Biochim Biophys Acta* **1548**, 23–28.
62. van Haaften RI, Haenen GR, Evelo CT, *et al.* (2002) Tocotrienols inhibit human glutathione S-transferase P1-1. *IUBMB Life* **54**, 81–84.
63. Mustacich DJ, Shields J, Horton RA, *et al.* (1998) Biliary secretion of  $\alpha$ -tocopherol and the role of the mdr2 P-glycoprotein in rats and mice. *Arch Biochem Biophys* **350**, 183–192.
64. Food and Drug Administration (2008) Consumer Health Information. <http://www.fda.gov/downloads/ForConsumers/ConsumerUpdates/ucm096391.pdf> (accessed July 2014).
65. Harper M & Poole A (2010) Diverse functions of protein kinase C isoforms in platelet activation and thrombus formation. *J Thromb Haemost* **8**, 454–462.
66. Ali M, Gudbranson CG & McDonald JWD (1980) Inhibition of human platelet cyclooxygenase by  $\alpha$ -tocopherol. *Prostaglandins Med* **4**, 79–85.
67. Freedman JE, Farhat JH, Loscalzo J, *et al.* (1996)  $\alpha$ -Tocopherol inhibits aggregation of human platelets by a protein kinase C-dependent mechanism. *Circulation* **94**, 2434–2440.
68. Freedman JE & Keaney JF Jr (2001) Vitamin E inhibition of platelet aggregation is independent of antioxidant activity. *J Nutr* **131**, 374S–377S.
69. Liu M, Wallmon A, Olsson-Mortlock C, *et al.* (2003) Mixed tocopherols inhibit platelet aggregation in humans: potential mechanisms. *Am J Clin Nutr* **77**, 700–706.
70. Steiner M (1983) Effect of  $\alpha$ -tocopherol administration on platelet function in man. *Thromb Haemost* **49**, 73–77.
71. Vane JR & Botting RM (2003) The mechanism of action of aspirin. *Thromb Res* **110**, 255–258.
72. Liede KE, Haukka JK, Saxen LM, *et al.* (1998) Increased tendency towards gingival bleeding caused by joint effect of  $\alpha$ -tocopherol supplementation and acetylsalicylic acid. *Ann Med* **30**, 542–546.
73. Steiner M, Glantz M & Lekos A (1995) Vitamin E plus aspirin compared with aspirin alone in patients with transient ischemic attacks. *Am J Clin Nutr* **62**, Suppl., 1381S–1384S.
74. Suttie JW (1980) Mechanism of action of vitamin K: synthesis of  $\gamma$ -carboxyglutamic acid. *CRC Crit Rev Biochem* **8**, 191–223.
75. Hirsh J, Dalen JE, Anderson DR, *et al.* (2001) Oral anti-coagulants: mechanism of action, clinical effectiveness, and optimal therapeutic range. *Chest* **108**, Suppl., 231S–246S.
76. Corrigan J & Ulfers L (1981) Effect of vitamin E on prothrombin levels in warfarin-induced vitamin K deficiency. *Am J Clin Nutr* **34**, 1701–1705.
77. Kim JM & White RH (1996) Effect of vitamin E on the anti-coagulant response to warfarin. *Am J Cardiol* **77**, 545–546.
78. Booth SL, Golly I, Sackeck JM, *et al.* (2004) Effect of vitamin E supplementation on vitamin K status in adults with normal coagulation status. *Am J Clin Nutr* **80**, 143–148.
79. Widdershoven J, Van Munster P, De Abreu R, *et al.* (1987) Four methods compared for measuring des-carboxy-prothrombin (PIVKA-II). *Clin Chem* **33**, 2074–2078.
80. Edson KZ, Prasad B, Unadkat JD, *et al.* (2013) Cytochrome P<sub>450</sub>-dependent catabolism of vitamin K:  $\omega$ -hydroxylation catalyzed by human CYP4F2 and CYP4F11. *Biochemistry* **52**, 8276–8285.
81. Farley SM, Leonard SW, Taylor AW, *et al.* (2013)  $\omega$ -Hydroxylation of phylloquinone by CYP4F2 is not increased by  $\alpha$ -tocopherol. *Mol Nutr Food Res* **57**, 1785–1793.
82. Shiotani A, Kamada T & Haruma K (2008) Low-dose aspirin-induced gastrointestinal diseases: past, present, and future. *J Gastroenterol* **43**, 581–588.
83. Fisher B, Redmond C, Legault-Poisson S, *et al.* (1990) Post-operative chemotherapy and tamoxifen compared with tamoxifen alone in the treatment of positive-node breast cancer patients aged 50 years and older with tumors responsive to tamoxifen: results from the National Surgical Adjuvant Breast and Bowel Project B-16. *J Clin Oncol* **8**, 1005–1018.
84. Shou J, Massarweh S, Osborne CK, *et al.* (2004) Mechanisms of tamoxifen resistance: increased estrogen receptor-HER2/neu cross-talk in ER/HER2-positive breast cancer. *J Natl Cancer Inst* **96**, 926–935.
85. Chang H, Huang J, Wang J, *et al.* (2002) Tamoxifen-induced increases in cytoplasmic free Ca<sup>2+</sup> levels in human breast cancer cells. *Breast Cancer Res Treat* **71**, 125–131.
86. Peralta EA, Viegas ML, Louis S, *et al.* (2006) Effect of vitamin E on tamoxifen-treated breast cancer cells. *Surgery* **140**, 607–615.
87. Peralta EA, Brewer AT, Louis S, *et al.* (2009) Vitamin E increases biomarkers of estrogen stimulation when taken with tamoxifen. *J Surg Res* **153**, 143–147.
88. Baliga R, Ueda N, Walker PD, *et al.* (1997) Oxidant mechanisms in toxic acute renal failure. *Am J Kidney Dis* **29**, 465–477.
89. Blackhall ML, Fassett RG, Sharman JE, *et al.* (2005) Effects of antioxidant supplementation on blood cyclosporin A and glomerular filtration rate in renal transplant recipients. *Nephrol Dial Transplant* **20**, 1970–1975.
90. de Vries AP, Oterdoom LH, Gans RO, *et al.* (2006) Supplementation with anti-oxidants vitamin C and E decreases cyclosporine A trough-levels in renal transplant recipients. *Nephrol Dial Transplant* **21**, 231–232.
91. Lake KD, Aaronson KD, Gorman LE, *et al.* (2005) Effect of oral vitamin E and C therapy on calcineurin inhibitor levels in heart transplant recipients. *J Heart Lung Transplant* **24**, 990–994.
92. Barany P, Stenvinkel P, Ottosson-Seeberger A, *et al.* (2001) Effect of 6 weeks of vitamin E administration on renal haemodynamic alterations following a single dose of neoral in healthy volunteers. *Nephrol Dial Transplant* **16**, 580–584.
93. Hebert M (1997) Contributions of hepatic and intestinal metabolism and P-glycoprotein to cyclosporine and tacrolimus oral drug delivery. *Adv Drug Deliv Rev* **27**, 201–214.
94. Vacca JP & Condra JH (1997) Clinically effective HIV-1 protease inhibitors. *Drug Discov Today* **2**, 261–272.
95. Dressman J, Kincer J, Matveev SV, *et al.* (2003) HIV protease inhibitors promote atherosclerotic lesion formation independent of dyslipidemia by increasing CD36-dependent cholesterol ester accumulation in macrophages. *J Clin Invest* **111**, 389–397.
96. Munteanu A, Zingg JM, Ricciarelli R, *et al.* (2005) CD36 overexpression in ritonavir-treated THP-1 cells is reversed by  $\alpha$ -tocopherol. *Free Radic Biol Med* **38**, 1047–1056.
97. Barella L, Muller PY, Schlachter M, *et al.* (2004) Identification of hepatic molecular mechanisms of action of  $\alpha$ -tocopherol using global gene expression profile analysis in rats. *Biochim Biophys Acta* **1689**, 66–74.



98. Gaedicke S, Zhang X, Schmelzer C, *et al.* (2008) Vitamin E dependent microRNA regulation in rat liver. *FEBS Lett* **582**, 3542–3546.
99. Meydani SN, Meydani M, Rall L, *et al.* (1994) Assessment of the safety of high-dose, short-term supplementation with vitamin E in healthy older adults. *Am J Clin Nutr* **60**, 704–709.
100. Meydani M, Lipman R, Han S, *et al.* (1998) The effect of long-term dietary supplementation with antioxidants. *Ann N Y Acad Sci* **854**, 352–360.
101. Milman U, Blum S, Shapira C, *et al.* (2008) Vitamin E supplementation reduces cardiovascular events in a subgroup of middle-aged individuals with both type 2 diabetes mellitus and the haptoglobin 2–2 genotype: a prospective double-blinded clinical trial. *Arterioscler Thromb Vasc Biol* **28**, 341–347.
102. Vardi M, Blum S & Levy AP (2012) Haptoglobin genotype and cardiovascular outcomes in diabetes mellitus - natural history of the disease and the effect of vitamin E treatment. Meta-analysis of the medical literature. *Eur J Intern Med* **23**, 628–632.
103. Blum S, Vardi M, Brown JB, *et al.* (2010) Vitamin E reduces cardiovascular disease in individuals with diabetes mellitus and the haptoglobin 2–2 genotype. *Pharmacogenomics* **11**, 675–684.
104. Scientific Committee on Food (SCF) (2003) Opinion of the Scientific Committee on Food on the Tolerable Upper Intake Level of Vitamin E. [http://ec.europa.eu/food/fs/sc/scf/out195\\_en.pdf](http://ec.europa.eu/food/fs/sc/scf/out195_en.pdf)