

A low-luminosity type-1 QSO sample: Insight from integral-field spectroscopy

Gerold Busch¹, Semir Smajić^{1,2}, Lydia Moser¹, Mónica Valencia-S.¹,
 Jens Zuther¹, Julia Scharwächter³ and Andreas Eckart^{1,2}

¹I. Physikalisches Institut, Universität zu Köln, Cologne, Germany
 email: busch@ph1.uni-koeln.de

²Max Planck Institut für Radioastronomie, Bonn, Germany

³LERMA, Observatoire de Paris, Paris, France

Abstract. The properties of the host galaxies of quasi-stellar objects (QSOs) are essential for the understanding of the suspected coevolution of central supermassive black holes (BHs) and their host galaxies. In Busch *et al.* (2014), we present a study of 20 low-luminosity type-1 QSOs (LLQSO; Busch *et al.* 2012) that have been selected from the Hamburg/ESO survey for bright UV-excess QSOs ($z \leq 0.06$). Performing careful decomposition of deep near-infrared J,H,K images, we found that the observed sources do not follow published $M_{\text{BH}} - L_{\text{bulge}}$ relations for inactive galaxies, supporting similar results found for type-1 AGN in the optical. This can be explained by overluminous bulges with very young stellar populations or undermassive black holes that are observed in a phase of growth. We use 3d-spectroscopy in the optical and near-infrared as a powerful tool to analyze gas and stellar kinematics, determine gas masses, star formation rate, trace underlying stellar continuum, etc. (e.g., Smajić *et al.* 2014) and thereby constrain possible evolution scenarios. The results will be interpreted in the context of galaxy evolution and particularly the still unknown role of the AGN in this process. Here, we show first results for HE 1029-1831.

Keywords. galaxies: active - galaxies: Seyfert - galaxies: evolution




Figure 1. From left to right: (1) In the $M_{\text{BH}} - L_{\text{bulge}}$ relation, the observed LLQSOs clearly lie off set compared to inactive galaxies collected by Kormendy & Ho (2013) (see Busch *et al.* 2014). (2) K -band image of the LLQSO HE 1029-1831 from Fischer *et al.* (2006). (3) Pa α map of HE 1029-1831 with a FOV of 8×8 arcsec². (4) the same with FOV of 3×3 arcsec². The atomic gas is distributed in spiral arms and a clumpy circum-nuclear ring, indicating star formation. This motivates further investigation of the star formation properties in the centers of LLQSOs.

References

- Busch G., Zuther J., Valencia-S. M., *et al.*, 2014, *A&A*, 561, A140
 Busch G., Zuther J., Valencia-S. M., *et al.*, 2012, PoS(Seyfert2012)060, arXiv:1307.1049
 Fischer S., Iserlohe C., Zuther J., *et al.*, 2006, *A&A*, 452, 827
 Kormendy J. & Ho L., 2013, *ARA&A*, 51, 511
 Smajić S., Moser L., Eckart A., *et al.*, 2014, *A&A*, 567, A119