

A magnetically collimated jet from the evolved star W43A

W. H. T. Vlemmings¹, P. J. Diamond¹ and H. Imai²

¹Jodrell Bank Observatory, Univ. of Manchester, Macclesfield, Cheshire SK11 9DL, U.K.
email: wouter@jb.man.ac.uk

²Department of Physics, Faculty of Science, Kagoshima University, Kagoshima 890-0065, Japan

Abstract. We present the first direct measurements of the magnetic field strength and direction in a collimated jet from an evolved star on its way to become a planetary nebula. Very Long Baseline Array (VLBA) observations of the linear and circular polarization of the H₂O masers in the collimated jet of W43A reveal a strong toroidal magnetic field, indicating that the jet is magnetically collimated. The magnetic field strength in the jet extrapolated back to the stellar surface yields a surface field of several Gauss, consistent with the measurements of maser polarization in a large sample of evolved stars. The origin of the magnetic field is yet unknown, although the jet precession might point to the existence of a heavy planet or stellar companion. This is the first direct observational evidence for magnetic collimation in the jets, that likely plays an important role in shaping planetary nebulae.

Keywords. magnetic fields, masers, stars: AGB and post-AGB

1. Introduction

W43A is an evolved star at a distance of 2.6 kpc (Diamond *et al.* 1985). It is surrounded by a thick circumstellar envelope (CSE) that exhibits OH, H₂O and SiO masers (Imai *et al.* 2005 and references therein). The H₂O masers of W43A occur in two clusters at ~1000 AU from the star near the opposing tips of a collimated jet. The jet, with a velocity of 145 km s⁻¹, has an inclination of 39° with respect to the sky plane, and a position angle of 65°. It precesses by 5° with a period of 55 years. The dynamical age of the jet is inferred to be only approximately 50 years (Imai *et al.* 2002). W43A is interpreted as belonging to a class of objects undergoing a rapid transition from an evolved star into a planetary nebula (PNe). Currently, owing to their short expected lifetime of less than 1000 years, only 4 sources of this class have been identified (Imai *et al.* 2002, Morris *et al.* 2003, Imai *et al.* 2004, Boboltz & Marvel 2005).

Maser polarization observations have revealed the magnetic field throughout the CSEs of a large number of evolved stars. Close to the central star, at typically 2 stellar radii, SiO masers indicate ordered fields of the order of several Gauss (e.g. Kembell & Diamond 1997, Herpin *et al.* 2006). At the outer edge of the CSE, the polarization measurements of OH masers reveal milliGauss magnetic fields and indicate weak alignment with CSE structure (Etoka & Diamond 2004). Recently, Zeeman splitting measurements of H₂O masers in the CSEs of a sample of evolved stars revealed large scale magnetic fields with field strengths between a hundred milliGauss up to a few Gauss (Vlemmings *et al.* 2002, Vlemmings *et al.* 2005). While the origin of the magnetic field is still unclear, theoretical models have shown that a dynamo between the slowly rotating stellar outer layers and the faster rotating core can produce the observed magnetic fields (Blackman *et al.* 2001). However, the required additional source of angular momentum to maintain the magnetic




Figure 1. The H_2O masers in the precessing jet (dashed-dotted line) of W43A (indicated by the cross). (top): The maser features with the determined linear polarization vectors scaled linearly according to the fractional linear polarization. The polarization vectors lie predominantly along the jet with a median angle of $\chi = 63 \pm 12^\circ$ east of north. (bottom): The toroidal magnetic field of W43A. The vectors indicate the determined magnetic field direction, perpendicular to the polarization vectors, at the location of the H_2O masers. The ellipses indicate the toroidal field along the jet, scaled with magnetic field strength $\propto r^{-1}$.

field likely requires the presence of a binary companion or heavy planet (Blackman *et al.* 2004, Frank *et al.* 2004, Nordhaus & Blackman 2006).

Magnetic fields around evolved stars are thought to be one of the main factors in shaping the CSEs and producing the asymmetries during the evolution of a spherically symmetric star into the often asymmetric PNe (Frank 2006, this volume). Similar to well established theories of collimated outflows in young stellar objects, theoretical models show that magnetic fields could be the collimating agents of the bi-polar jets in young proto-planetary nebulae such as W43A (e.g. García-Segura *et al.* 2005). Here we describe the first direct detection of a magnetically collimated jet from the evolved star W43A, recently published by Vlemmings *et al.* (2006a) (hereafter V06a).




Figure 2. (left): The total intensity and circular polarization spectrum of the H₂O maser feature for which circular polarization was detected. (right): The figure, reproduced from Vlemmings *et al.* (2005), of measured magnetic fields for the masers in the CSEs of evolved stars. The dashed-dotted boxes indicate the range of magnetic field strengths measured for the SiO and OH masers of Mira stars and the solid boxes those of supergiant stars. The thin arrows indicate the H₂O maser magnetic fields measured by Vlemmings *et al.* (2002) and Vlemmings *et al.* (2005), where the lengths of the arrows indicate the thickness of the H₂O maser shell with the symbols drawn on the outer edge. The symbols without arrows are the measurements on SiO and OH masers from the literature on the same sample of stars. The thick solid arrow indicates the magnetic field in the jet of W43A.

2. Observations

The observations of W43A were performed with the NRAO[†] VLBA on December 8 2004 at the frequency of the 6₁₆–5₂₃ rotational transition of H₂O, 22.235080 GHz. The correlation and data-reduction was performed as described by Vlemmings *et al.* (2006b), where similar methods were used to study H₂O maser magnetic fields in a high-mass star-forming region. Polarization calibration was performed with respect to the calibrator J1743-0350 and we estimate the systematic error of the polarization angles to be at most $\sim 8^\circ$.

3. Results

Linear polarizations were determined for several 22 GHz H₂O maser features in both the red-shifted and blue-shifted tip of the jet of W43A. The linear polarization vectors are either parallel or perpendicular to the magnetic field (Goldreich *et al.* 1973). As discussed in V06a and elaborated upon in Vlemmings *et al.* (in preparation), we find that the polarization vectors of the masers of W43A are mostly perpendicular to the magnetic field direction. The observed maser features and their linear polarization vectors are shown at the top of Fig.1 and the derived magnetic field direction is shown in the

[†] The National Radio Astronomy Observatory (NRAO) is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

bottom panel of the figure. In addition to the linear polarization, we detected circular polarization of $P_V = 0.33 \pm 0.09\%$ (Fig. 2, left), which as shown by V06a indicates a deprojected magnetic field strength of $B = 200 \pm 78$ mG.

4. Discussion

As the H₂O masers are excited in swept up material in the jet, the magnetic field in the maser region is enhanced. Partial coupling of the magnetic field to the gas indicates the toroidal magnetic field strength around the collimated jet, at ~ 1000 AU from W43A is $B_\phi \sim 2$ mG (V06a). The linear polarization vectors, perpendicular to the jet axis, indicate that we are tracing a confining toroidal magnetic field. A toroidal field depends on distance from the star r as $B_\phi \propto r^{-1}$ and we thus find that at the surface of the star, the magnetic field strength is between ~ 2 – 20 G. As shown in Fig. 2 (right) this is fully consistent with earlier measurements of the magnetic field in the envelopes of evolved stars. SiO maser observations (Imai et al. 2005) seem to indicate that the jet formation occurs within the SiO maser region, close to the star, and thus that the collimating agent is related to the star itself. Considering the shape and strength of the magnetic field in the jet of W43A, we conclude that the magnetic field is the primary collimating agent, making this the first direct detection of magnetic collimation in any astrophysical jet.

Acknowledgements

The work by WV has been supported by a Marie Curie Intra-European fellowship within the 6th European Community Framework Program under contract number MEIF-CT-2005-010393.

References

- Blackman, E. G. 2004, in *ASP Conf. Ser. 313: Asymmetrical Planetary Nebulae III: Winds, Structure and the Thunderbird*, 401
- Blackman, E. G., Frank, A., Markiel, J. A., Thomas, J. H., & Van Horn, H. M. 2001, *Nature*, 409, 485
- Boboltz, D. A. & Marvel, K. B. 2005, *ApJ*, 627, L45
- Diamond, P. J., Norris, R. P., Rowland, P. R., Booth, R. S., & Nyman, L.-A. 1985, *MNRAS*, 212, 1
- Etoka, S. & Diamond, P. 2004, *MNRAS*, 348, 34
- Frank, A. & Blackman, E. G. 2004, *ApJ*, 614, 737
- García-Segura, G., López, J. A., & Franco, J. 2005, *ApJ*, 618, 919
- Goldreich, P., Keeley, D. A., & Kwan, J. Y. 1973, *ApJ*, 179, 111
- Herpin, F., Baudry, A., Thum, C., Morris, D., & Wiesemeyer, H. 2006, *A&A*, 450, 667
- Imai, H., Morris, M., Sahai, R., Hachisuka, K., & Azzollini F., J. R. 2004, *A&A*, 420, 265
- Imai, H., Nakashima, J.-i., Diamond, P. J., Miyazaki, A., & Deguchi, S. 2005, *ApJ*, 622, L125
- Imai, H., Obara, K., Diamond, P. J., Omodaka, T., & Sasao, T. 2002, *Nature*, 417, 829
- Kemball, A. J. & Diamond, P. J. 1997, *ApJ*, 481, L111
- Morris, M. R., Sahai, R., & Claussen, M. 2003, in *Rev. Mexicana AyA Conference Series*, 20–22
- Nordhaus, J., & Blackman, E.G. 2006, *MNRAS*, 370, 2004
- Vlemmings, W. H. T., Diamond, P. J., & Imai, H. 2006a, *Nature*, 440, 58
- Vlemmings, W. H. T., Diamond, P. J., & van Langevelde, H. J. 2002, *A&A*, 394, 589
- Vlemmings, W. H. T., Diamond, P. J., van Langevelde, H. J., & Torrelles, J. M. 2006b, *A&A*, 448, 597
- Vlemmings, W. H. T., van Langevelde, H. J., & Diamond, P. J. 2005, *A&A*, 434, 1029