

ROOTS OF UNITY AS QUOTIENTS OF TWO ROOTS OF A POLYNOMIAL

ARTŪRAS DUBICKAS

(Received 1 March 2011; accepted 1 February 2012)

Communicated by I. E. Shparlinski

Dedicated to the memory of Alf van der Poorten

Abstract

Let K be a number field. For $f \in K[x]$, we give an upper bound on the least positive integer $T = T(f)$ such that no quotient of two distinct T th powers of roots of f is a root of unity. For each $\varepsilon > 0$ and each $f \in \mathbb{Q}[x]$ of degree $d \geq d(\varepsilon)$ we prove that $\log T(f) < (2 + \varepsilon)\sqrt{d \log d}$. In the opposite direction, we show that the constant 2 cannot be replaced by a number smaller than 1. These estimates are useful in the study of degenerate and nondegenerate linear recurrence sequences over a number field K .

2010 *Mathematics subject classification*: primary 11R04; secondary 11R18, 11B37.

Keywords and phrases: root of unity, number field, linear recurrence sequence.

1. Introduction

Let K be a number field, that is, a finite extension of the field of rational numbers \mathbb{Q} , and let

$$f(x) := x^d + a_{d-1}x^{d-1} + \cdots + a_0 = \prod_{i=1}^d (x - \alpha_i) \in K[x]$$

be a monic polynomial of degree $d \geq 2$. It may happen that the quotient of two distinct roots of f , say, α_i/α_j , $1 \leq i < j \leq d$, is a root of unity (for example, for $f(x) = x^d - 2 \in \mathbb{Q}[x]$). We say that the polynomial f is *torsion-free* if no quotient of two distinct roots of f is a root of unity. In the case where $f(x) = \prod_{i=1}^d (x - \alpha_i)$ is not torsion-free, the polynomial $f_T(x) := \prod_{i=1}^d (x - \alpha_i^T)$ is torsion-free for some $T \in \mathbb{N}$. The smallest positive integer $T = T(f)$ with this property is called the *torsion order* of f , so that torsion-free polynomials have torsion order one. For example, for the polynomial $x^d - 2$, we have $T(x^d - 2) = d$. The torsion order of a polynomial can be greater than its degree, for example, $T(\Phi_d) = d$, where Φ_d is the d th cyclotomic polynomial of degree $\varphi(d)$. Note that the torsion order $T(f)$ is independent of the field K . If $f \in K[x]$

is not monic and has a leading coefficient $a \in K \setminus \{0\}$ then the torsion order of f is defined as $T(f/a)$.

Here is the main result of this paper.

THEOREM 1. *Let K be a number field with $k := [K : \mathbb{Q}]$. For every $f \in K[x]$ of degree $d \geq 2$,*

$$\log T(f) < (1.053\,14 + \sqrt{6k})\sqrt{d \log(kd)}. \tag{1}$$

Furthermore, if $\varepsilon > 0$ then

$$\log T(f) < (1 + \varepsilon)(1 + \sqrt{k})\sqrt{d \log d} \tag{2}$$

provided that d is large enough. On the other hand, for all $\varepsilon > 0$ there exists $d_0(\varepsilon)$ such that for each positive integer $d \geq d_0(\varepsilon)$ there is a monic polynomial $\Phi \in \mathbb{Z}[x] \subset K[x]$ of degree d for which

$$\log T(\Phi) > (1 - \varepsilon)\sqrt{d \log d}. \tag{3}$$

On replacing ε by $\varepsilon/2$ in (2) and selecting $K := \mathbb{Q}$, we find that

$$\exp((1 - \varepsilon)\sqrt{d \log d}) < \max_{f \in \mathbb{Q}[x], \deg f = d} T(f) < \exp((2 + \varepsilon)\sqrt{d \log d}) \tag{4}$$

for each $\varepsilon > 0$ and each sufficiently large d . Note that the difference between the upper and lower bounds in (4) is only in the constants 1 and 2.

The torsion order T of the polynomial $f(x) = x^d + a_{d-1}x^{d-1} + \dots + a_0$ is related to the properties of the linear recurrence sequence

$$x_{n+d} = -a_{d-1}x_{n+d-1} - \dots - a_0x_n,$$

where $n = 1, 2, \dots$. Such a sequence is called *degenerate* if its characteristic polynomial f has a pair of distinct roots whose ratio is a root of unity and *nondegenerate* otherwise. On replacing f by f_T , where T is the torsion order of f , one reduces the study of arbitrary linear recurrence sequences to the study of nondegenerate sequences. Theorem 1.2 in [4] asserts that, for some $t \leq T(f)$, each subsequence x_{m+s} , where $n = 1, 2, \dots$, is either identically zero or nondegenerate. Other applications of the torsion order to the Skolem–Mahler–Lech theorem were mentioned by Berstel and Mignotte [2] (see also [4, Ch. 2]). Schinzel [11] used $T(f)$ to treat an old problem of Pólya [9] on the description of rational functions $\sum_{n=0}^{\infty} u_n x^n \in \mathbb{Q}(x)$ whose numerators are divisible by only finitely many primes. In this sense the bound in (4) gives the best possible (up to a constant) estimate in [11, Theorem 2]. See also [14], where the case $K = \mathbb{Q}$ was considered. Robba [10] investigated the case of a number field, but his upper bound $T(f) \leq 2^{kd+1}$ is weaker than that given in (1).

It seems likely that if $f \in \mathbb{Q}[x]$ is irreducible over the field \mathbb{Q} then the upper bound for $T(f)$ should be given by the inequality

$$\varphi(T(f)) \leq d, \tag{5}$$

where φ is Euler’s function. Note that equality holds in (5) for cyclotomic polynomials $f = \Phi_d$. Since $\liminf_{n \rightarrow \infty} (\varphi(n) \log \log n)/n = e^{-\gamma}$, where $\gamma = 0.577\,21\dots$ is Euler’s

constant (see, for example, [5, Theorem 324]), the upper bound (5) via [11, Lemma 3] would imply the inequality

$$T(f) < e^\gamma d(\log \log d + 4) \tag{6}$$

for every $d \geq 2$. Unfortunately, the proof of Corollary 2.1 (which is equivalent to (5)) in our paper [3] contains an error. The best known bound in this direction is due to Schinzel [11] who showed that

$$T(f) < e^{3\gamma/2} d^{3/2} (\log \log d + 4)^{3/2}$$

for every irreducible polynomial $f \in \mathbb{Q}[x]$ of degree $d \geq 2$ (which is short of the conjectured bound (6)). In [1, 3, 6, 14], one can find different proofs of the inequality

$$[K(\zeta) : K] \leq d = [K(\alpha) : K]$$

when α, α' are two algebraic numbers conjugate over K and $\zeta := \alpha/\alpha'$ is a root of unity. This shows that one can get rid of at least one root of unity among the ratios of roots of an irreducible polynomial $f \in K[x]$ of degree d by taking some power t for which $\varphi(t) \leq d$.

In the next section we shall prove (3) and give another example which shows that for some $f \in K[x]$ the torsion degree $T(f)$ may tend to infinity as $k \rightarrow \infty$ for cyclotomic extensions K of \mathbb{Q} . In Section 3 we state two results on the least common multiple of positive integers b_1, b_2, \dots, b_m , whose sum (or the sum of the values of Euler's function $\varphi(b_1), \varphi(b_2), \dots, \varphi(b_m)$) does not exceed some fixed integer n . We give two more lemmas in Section 4, and then complete the proofs of (1) and (2) in Section 5.

2. Examples

Let $X(d)$ be the largest integer for which

$$\sum_{p \leq X(d)} (p - 1) \leq d, \tag{7}$$

where the sum is taken over prime numbers p . Set

$$\Phi(x) := g(x) \prod_{p \leq X(d)} (x^{p-1} + \dots + x + 1), \tag{8}$$

where $g(x) := \prod_{i=1}^e (x - i) \in \mathbb{Z}[x]$ is a polynomial of degree

$$e := d - \sum_{p \leq X(d)} (p - 1).$$

Then $\Phi(x) \in \mathbb{Z}[x]$ is a monic polynomial of degree d . (Note that $\Phi(x) \in \mathbb{Z}[x] \subset K[x]$ for every number field K .) Put

$$Y(d) := \prod_{p \leq X(d)} p, \tag{9}$$

where the product is taken over primes p . Clearly, Φ_T is a torsion-free polynomial for $T = Y(d)$, since the T th powers of the roots of Φ are i^T , where $i = 1, \dots, e$, and 1, because $\zeta_p^{jY(d)} = 1$ whenever $p \leq X(d)$ and $j = 1, \dots, p - 1$. (Throughout, $\zeta_r := e^{2\pi i/r}$ is the primitive r th root of unity.) On the other hand, if some prime $p \leq X(d)$ does not divide T then Φ_T is not torsion-free, because the quotient of two distinct roots of Φ_T , say ζ_p^{2T} and ζ_p^T , is a root of unity. Consequently, for the polynomial defined in (8),

$$T(\Phi) = Y(d).$$

Now, using the prime number theorem $p_i \sim i \log i$ as $i \rightarrow \infty$, from (7) and (9) one can easily derive that

$$X(d) \sim \sqrt{d \log d} \quad \text{and} \quad \log Y(d) \sim X(d)$$

as $d \rightarrow \infty$. Hence the inequality $\log T(\Phi) = \log Y(d) > (1 - \varepsilon)\sqrt{d \log d}$ holds for every $d \geq d_0(\varepsilon)$ and every $\varepsilon > 0$. This proves (3).

Suppose next that $K = \mathbb{Q}(\zeta_m)$, where $\varphi(m) = k$, so that $k = [K : \mathbb{Q}]$. Let us take

$$G(x) := g_1(x) \prod_{m < p \leq X_1(d)} (x^p - \zeta_m),$$

where the product is taken over primes p , the number $X_1(d)$ is the largest integer for which $\sum_{m < p \leq X_1(d)} p \leq d$, and $g_1(x) := \prod_{i=1}^{e_1} (x - i)$ with $e_1 := d - \sum_{m < p \leq X_1(d)} p$. By the same argument as above, we see that $G \in K[x]$ is a monic polynomial of degree d whose torsion order is

$$T(G) = m \prod_{m < p \leq X_1(d)} p > m \exp((1 - \varepsilon)\sqrt{d \log d})$$

for each $\varepsilon > 0$ and each sufficiently large d . Since $m > k$, it follows that

$$\log T(G) > \log k + (1 - \varepsilon)\sqrt{d \log d}$$

for each $\varepsilon > 0$ and each sufficiently large d .

3. Two results for the least common multiple

LEMMA 2. *Let b_1, \dots, b_m be some positive integers satisfying $b_1 + \dots + b_m \leq n$, where $n \geq 2$. Then*

$$\text{lcm}(b_1, \dots, b_m) < \exp(1.053 \, 14 \sqrt{n \log n}),$$

where lcm stands for the least common multiple. Furthermore, for each $\varepsilon > 0$ there is an integer $n_0(\varepsilon)$ such that, for every $n \geq n_0(\varepsilon)$,

$$\text{lcm}(b_1, \dots, b_m) < \exp((1 + \varepsilon)\sqrt{n \log n}).$$

The first part of Lemma 2 was proved by Massias [7]. In fact, $\text{lcm}(b_1, \dots, b_m)$ is the order of an element consisting of m disjoint cycles of lengths b_1, \dots, b_m in the

full symmetric group on n elements S_n . Let $M(n)$ be the maximal order of an element in S_n . Landau proved that $\log M(n) \sim \sqrt{n \log n}$ as $n \rightarrow \infty$ and Shah [12] gave a more precise asymptotic formula

$$\log M(n) = \sqrt{n \log n} (1 + \log \log n / (2 \log n) + O(1/\log n)).$$

This implies the second part of Lemma 2. See also [8, 13] for further work on this problem. It is interesting to note that the upper bound $1.053\ 13\ \dots$ is attained for the symmetric group $S_{1\ 319\ 766}$.

The following lemma was proved by Berstel and Mignotte [2].

LEMMA 3. *Let b_1, \dots, b_m be positive integers satisfying $\varphi(b_1) + \dots + \varphi(b_m) \leq n$, where $n \geq 2$. Then*

$$\text{lcm}(b_1, \dots, b_m) < \exp(\sqrt{6n \log n}).$$

Furthermore, the constant $\sqrt{6}$ can be replaced by any constant strictly greater than 1 if n is sufficiently large.

4. Some divisibility results for the torsion order of a polynomial

Let $f \in K[x]$ be a separable polynomial of degree $d \geq 2$. We say that two of its roots α and α' belong to the same equivalence class if their quotient is a root of unity. Suppose that there are $s := s(f)$ distinct equivalence classes. It is easy to see that each equivalence class contains the same number of elements, say, $\ell := \ell(f)$ roots of f , so

$$d = s\ell = s(f)\ell(f). \tag{10}$$

Now, let $f \in K[x]$ be an irreducible polynomial of degree $d \geq 2$, and let α be one of its roots. Suppose that r is the largest positive integer for which

$$\mathbb{Q}(\zeta_r) \subseteq K(\alpha). \tag{11}$$

Then $K(\zeta_r) \subseteq K(\alpha)$. Since $K(\zeta_r)$ is a normal extension of K , the field $K(\zeta_r)$ is contained in the field $K(\alpha')$ for any conjugate α' of α over K . Hence $\mathbb{Q}(\zeta_r)$ is contained in every field $K(\alpha')$ too. Thus $r := r(f, K)$ is independent of the choice of α and depends only on the polynomial f and the field K . Of course,

$$r(f, K) = w(K(\alpha)), \tag{12}$$

where $w(F)$ stands for the number of distinct roots of unity lying in the field F .

We claim that if $\alpha \neq \alpha'$ and α/α' is a root of unity (so that $\ell \geq 2$) then

$$\alpha^{\ell r} = (\alpha')^{\ell r}. \tag{13}$$

Indeed, write $\gamma(C)$ for the product of conjugates of α belonging to the equivalence class C . From $\alpha, \alpha' \in C$ we obtain $\alpha^\ell = \zeta\gamma(C)$ and $(\alpha')^\ell = \zeta'\gamma(C)$ for some roots of unity ζ, ζ' . It is clear that $s = s(f)$ equivalence classes are blocks of imprimitivity of the Galois group $\text{Gal}(K(\alpha)/K)$. Hence every automorphism $\sigma \in \text{Gal}(K(\alpha)/K)$ satisfying $\sigma(\alpha) = \alpha$ is a permutation of the set C , so it maps $\gamma(C)$ to $\gamma(C)$. Thus $\gamma(C) \in K(\alpha)$.

This yields $\zeta = \alpha^\ell / \gamma(C) \in K(\alpha)$, so that $\zeta \in \mathbb{Q}(\zeta_r)$, by (11). Hence $\zeta^r = 1$. By the same argument, $(\zeta')^r = 1$. Therefore,

$$\left(\frac{\alpha}{\alpha'}\right)^{\ell r} = \left(\frac{\zeta \gamma(C)}{\zeta' \gamma(C)}\right)^r = \frac{\zeta^r}{(\zeta')^r} = 1.$$

This proves (13) and so implies the following result.

LEMMA 4. *The torsion order of an irreducible polynomial $f \in K[x]$ satisfies*

$$T(f) \mid \ell(f)r,$$

where $r := r(f, K)$ is defined in (11) and (12).

Next we extend this result to the product of two irreducible polynomials.

LEMMA 5. *Let $f_1, f_2 \in K[x]$ be two distinct irreducible polynomials having $\ell_1 := \ell(f_1)$ and $\ell_2 := \ell(f_2)$ elements in their equivalence classes and with $r_1 := r(f_1, K)$ and $r_2 := r(f_2, K)$ defined as in (11) and (12). Then*

$$T(f_1 f_2) \mid \text{lcm}(\ell_1 \text{lcm}(r_1, r_2), \ell_2 \text{lcm}(r_1, r_2)).$$

PROOF. In view of Lemma 4 it suffices to prove that if α is a root of f_1 and β is a root of f_2 such that α/β is a root of unity then

$$\alpha^L = \beta^L,$$

where $L := \text{lcm}(\ell_1 r, \ell_2 r)$ and $r := \text{lcm}(r_1, r_2)$. Suppose that α and β belong to the equivalence classes C_1 and C_2 . As above, let $\gamma(C_1)$ and $\gamma(C_2)$ be the product of conjugates of α from C_1 and the product of conjugates of β from C_2 . Then

$$\alpha^{\ell_1} = \xi_1 \gamma(C_1) \quad \text{and} \quad \beta^{\ell_2} = \xi_2 \gamma(C_2) \tag{14}$$

with some roots of unity ξ_1, ξ_2 .

Let S be the set of $\deg f_1 + \deg f_2$ roots of the polynomial $f_1 f_2$. The equivalence class of S containing α consists of $C_1 \cup C_2$. Consider the Galois group $\text{Gal}(K(\alpha, \beta)/K)$ as permutations of the set S . Each element $\sigma \in \text{Gal}(K(\alpha, \beta)/K)$ that fixes α (or β) permutes the class C_1 and the class C_2 . Hence the products $\gamma(C_1)$ and $\gamma(C_2)$ both lie in the intersection $K(\alpha) \cap K(\beta)$. Thus $\xi_1 \in K(\alpha)$ and $\xi_2 \in K(\beta)$, by (14). From (14), the quotient $\zeta := \gamma(C_1)^{L/\ell_1 r} / \gamma(C_2)^{L/\ell_2 r}$ is a root of unity, because α/β is a root of unity. Note that $\zeta, \xi_1 \in K(\alpha)$ yields $\zeta, \xi_1 \in \mathbb{Q}(\zeta_{r_1})$, by the definition of r_1 . Similarly, $\zeta, \xi_2 \in K(\beta)$ yields $\zeta, \xi_2 \in \mathbb{Q}(\zeta_{r_2})$. It follows that ζ, ξ_1, ξ_2 lie in the compositum

$$\mathbb{Q}(\zeta_{r_1})\mathbb{Q}(\zeta_{r_2}) = \mathbb{Q}(\zeta_{r_1, r_2}) = \mathbb{Q}(\zeta_{\text{lcm}(r_1, r_2)}) = \mathbb{Q}(\zeta_r).$$

Now, using the fact that the root of unity $\zeta \xi_1^{L/\ell_1 r} \xi_2^{-L/\ell_2 r}$ belongs to the field $\mathbb{Q}(\zeta_r)$ (so that its r th power is 1), we obtain

$$\frac{\alpha^L}{\beta^L} = \frac{\xi_1^{L/\ell_1} \gamma(C_1)^{L/\ell_1}}{\xi_2^{L/\ell_2} \gamma(C_2)^{L/\ell_2}} = \left(\frac{\xi_1^{L/\ell_1 r} \zeta}{\xi_2^{L/\ell_2 r}}\right)^r = 1,$$

which completes the proof of the lemma. □

5. Proof of the upper bound for torsion order

Without loss of generality, we may assume that $f \in K[x]$ is monic. Let us write f in the form

$$f(x) = f_1(x)^{k_1} f_2(x)^{k_2} \cdots f_m(x)^{k_m}$$

for distinct irreducible monic polynomials $f_1, \dots, f_m \in K[x]$ and $k_1, \dots, k_m \in \mathbb{N}$. Obviously, $T(f) = T(f_1 \cdots f_m)$ and the degree of the polynomial $f_1 \cdots f_m$ is smaller than $\deg f$ provided that at least one k_i is greater than 1. So we may assume that $k_1 = \cdots = k_m = 1$. Then, by Lemmas 4 and 5, for $f = f_1 \cdots f_m$,

$$T(f) \mid \text{lcm}(\ell_1, \dots, \ell_m) \text{lcm}(r_1, \dots, r_m) \tag{15}$$

with $r_i := r(f_i, K)$ defined in (11) and (12), because $\text{lcm}(\ell_i \text{lcm}(r_i, r_j), \ell_j \text{lcm}(r_i, r_j))$ divides $\text{lcm}(\ell_i, \ell_j) \text{lcm}(r_i, r_j)$ for all indices i, j in the range $1 \leq i, j \leq m$. From (10) we see that $\ell_i \mid d_i$, where $d_i = \deg f_i$. Thus $\text{lcm}(\ell_1, \dots, \ell_m)$ divides $\text{lcm}(d_1, \dots, d_m)$. Using the equality $\sum_{i=1}^m d_i = d = \deg f$ from Lemma 2 we deduce that

$$\text{lcm}(\ell_1, \dots, \ell_m) \leq \text{lcm}(d_1, \dots, d_m) < \exp(1.053 \, 14 \sqrt{d \log d}). \tag{16}$$

Suppose that α_i is a root of f_i for $i = 1, \dots, m$. Then we see that $\mathbb{Q}(\zeta_{r_i}) \subseteq K(\alpha_i)$ from (11). Hence $[\mathbb{Q}(\zeta_{r_i}) : \mathbb{Q}] = \varphi(r_i)$ divides

$$[K(\alpha_i) : \mathbb{Q}] = [K : \mathbb{Q}] \cdot [K(\alpha_i) : K] = kd_i.$$

In particular, $\varphi(r_i) \leq kd_i$ for each $i = 1, \dots, m$ and thus

$$\sum_{i=1}^m \varphi(r_i) \leq k \sum_{i=1}^m d_i = kd.$$

Therefore, Lemma 3 implies that

$$\text{lcm}(r_1, \dots, r_m) < \exp(\sqrt{6kd \log(kd)}). \tag{17}$$

Now, from (15), multiplying (16) and (17) we derive (1).

Next, suppose that $\varepsilon > 0$ and that d is large enough. Then, by Lemmas 2 and 3, inequality (16) becomes

$$\text{lcm}(\ell_1, \dots, \ell_m) \leq \text{lcm}(d_1, \dots, d_m) < \exp((1 + \varepsilon)\sqrt{d \log d}),$$

whereas (17) is replaced by

$$\text{lcm}(r_1, \dots, r_m) < \exp((1 + \varepsilon/2)\sqrt{kd \log(kd)}) < \exp((1 + \varepsilon)\sqrt{kd \log d}).$$

Multiplying these two inequalities, we derive (2) in view of (15).

References

- [1] M. G. Aschbacher and R. M. Guralnick, ‘On Abelian quotients of primitive groups’, *Proc. Amer. Math. Soc.* **107** (1989), 89–95.
- [2] J. Berstel and M. Mignotte, ‘Deux propriétés décidables des suites récurrentes linéaires’, *Bull. Soc. Math. France* **104** (1976), 175–194.
- [3] P. Drungilas and A. Dubickas, ‘On subfields of a field generated by two conjugate algebraic numbers’, *Proc. Edinb. Math. Soc.* **47** (2004), 119–123.
- [4] G. Everest, A. van der Poorten, I. Shparlinski and T. Ward, *Recurrence Sequences*, Mathematical Surveys and Monographs, 104 (American Mathematical Society, Providence, RI, 2003).
- [5] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers* (Oxford University Press, Oxford, 1979).
- [6] I. M. Isaacs, ‘Quotients which are roots of unity (solution of problem 6523)’, *Amer. Math. Monthly* **95** (1988), 561–562.
- [7] J.-P. Massias, ‘Majoration explicite de l’ordre maximum d’un élément du groupe symétrique’, *Ann. Fac. Sci. Toulouse Math.* **6** (1984), 269–281.
- [8] J.-P. Massias, J.-L. Nicolas and G. Robin, ‘Evaluation asymptotique de l’ordre maximum d’un élément du groupe symétrique’, *Acta Arith.* **104** (1988), 221–242.
- [9] G. Pólya, ‘Arithmetische Eigenschaften der Reihenentwicklungen rationaler Funktionen’, *J. reine angew. Math.* **151** (1921), 1–31.
- [10] Ph. Robba, ‘Zéros de suites récurrentes linéaires’, *Groupe Étude Anal. Ultramétrie*, 5 (1977/78), Exposé No. 13, 1978, 5 pp.
- [11] A. Schinzel, ‘Around Pólya’s theorem on the set of prime divisors of a linear recurrence’, in: *Diophantine Equations. Papers from the international conference held in honor of T. N. Shorey’s 60th birthday*, Mumbai, 16–20 December 2005, (ed. N. Saradha) (Narosa Publishing House, New Delhi, 2008), pp. 225–233.
- [12] S. Shah, ‘An inequality for the arithmetical function $g(x)$ ’, *J. Indian Math. Soc.* **3** (1939), 316–318.
- [13] M. Szalay, ‘On the maximal order in S_n and S_n^{**} ’, *Acta Arith.* **37** (1980), 321–331.
- [14] K. Yokoyama, Z. Li and I. Nemes, ‘Finding roots of unity among quotients of the roots of an integral polynomial’, in: *Proceedings of the 1995 International Symposium on Symbolic and Algebraic Computation, ISSAC’95*, Montreal, 10–12 July 1995, (ed. A. H. M. Levelt) (ACM Press, New York, 1995), pp. 85–89.

ARTŪRAS DUBICKAS, Department of Mathematics and Informatics,
 Vilnius University, Naugarduko 24, Vilnius LT-03225, Lithuania
 e-mail: arturas.dubickas@mif.vu.lt