

## Original Article

**Cite this article:** van der Velden NCA, Smets EMA, van Vliet LM, Brom L, van Laarhoven HWM, Henselmans I (2024) Effects of prognostic communication strategies on emotions, coping, and appreciation of consultations: An experimental study in advanced cancer. *Palliative and Supportive Care*, 1–13. <https://doi.org/10.1017/S1478951524000403>

Received: 4 August 2023  
Revised: 15 February 2024  
Accepted: 19 February 2024


**Keywords:**

Prognosis; neoplasms; communication; physician–patient relations; emotions; patient satisfaction

**Corresponding author:**

Naomi C.A. van der Velden;  
Email: [n.c.vandervelden@amsterdamumc.nl](mailto:n.c.vandervelden@amsterdamumc.nl)

# Effects of prognostic communication strategies on emotions, coping, and appreciation of consultations: An experimental study in advanced cancer

Naomi C.A. van der Velden, <sup>PH.D.</sup>1,2,3 , Ellen M.A. Smets, <sup>PH.D.</sup>1,2,3, Liesbeth M. van Vliet, <sup>PH.D.</sup>4, Linda Brom, <sup>PH.D.</sup>5,6, Hanneke W.M. van Laarhoven, <sup>M.D., PH.D.</sup>3,7 and Inge Henselmans, <sup>PH.D.</sup>1,2,3

<sup>1</sup>Department of Medical Psychology, Amsterdam UMC Location University of Amsterdam, Amsterdam, The Netherlands; <sup>2</sup>Quality of Care, Amsterdam Public Health, Amsterdam, The Netherlands; <sup>3</sup>Cancer Treatment and Quality of Life, Cancer Center Amsterdam, Amsterdam, The Netherlands; <sup>4</sup>Department of Health, Medical and Neuropsychology, University of Leiden, Leiden, The Netherlands; <sup>5</sup>Department of Research and Development, Netherlands Comprehensive Cancer Organisation (IKNL), Utrecht, The Netherlands; <sup>6</sup>Netherlands Association for Palliative Care (PZNL), Utrecht, The Netherlands and <sup>7</sup>Department of Medical Oncology, Amsterdam UMC Location University of Amsterdam, Amsterdam, The Netherlands

**Abstract**

**Objectives.** We aimed to investigate effects of prognostic communication strategies on emotions, coping, and appreciation of consultations in advanced cancer.

**Methods.** For this experimental study, we created 8 videos of a scripted oncological consultation, only varying in prognostic communication strategies. Disease-naïve individuals ( $n = 1036$ ) completed surveys before and after watching 1 video, while imagining being the depicted cancer patient. We investigated effects of the *type* of disclosure (prognostic disclosure vs. communication of unpredictability vs. non-disclosure) and *content* of disclosure (standard vs. standard and best-case vs. standard, best- and worst-case survival scenarios; numerical vs. word-based estimates) on emotions, coping, and appreciation of consultations. Moderating effects of individual characteristics were tested.

**Results.** Participants generally reported more satisfaction ( $p < .001$ ) after prognostic disclosure versus communication of unpredictability and less uncertainty ( $p = .042$ ), more satisfaction ( $p = .005$ ), and more desirability ( $p = .016$ ) regarding prognostic information after numerical versus word-based estimates. Effects of different survival scenarios were absent. Prognostic communication strategies lacked effects on emotions and coping. Significant moderators included prognostic information preference and uncertainty tolerance.

**Significance of results.** In an experimental setting, prognostic disclosure does not cause more negative emotions than non-disclosure and numerical estimates are more strongly appreciated than words. Oncologists' worries about harming patients should not preclude disclosing (precise) prognostic information, yet sensitivity to individual preferences and characteristics remains pivotal.

**Introduction**

Patients who are diagnosed with advanced cancer are confronted with an uncertain future (Shilling et al. 2017). Information about prognosis helps patients to adjust to the last phase of life emotionally (Butow et al. 2020; van der Velden et al. 2020). The majority of patients with advanced cancer want prognostic information (Innes and Payne 2009; van der Velden et al. 2022b). Those who prefer not knowing often want to stay optimistic and foster hope (van der Velden et al. 2022b).

Prognostication is difficult, as it is hard to predict when individual patients will die from their cancer and group-level survival data may be limited. While oncologists recognize the importance of prognostic information (Liu et al. 2014), they regularly report discomfort with disclosing prognosis (Butow et al. 2020). They worry about undermining patients' hope and fighting spirit or may not know how to respond to the expected raise in negative emotions, such as anxiety (Hancock et al. 2007). Oncologists furthermore fear that prognostic disclosure harms patients' trust in them, patients' appreciation of the consultation, or the therapeutic relationship (Daugherty and Hlubocky 2008; Gordon and Daugherty 2003; Sisk and Mack 2018).

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

According to observational studies, oncologists often explicate the unpredictability of prognosis or forgo prognostic discussions altogether (Chou *et al.* 2017; Derry *et al.* 2019a; Gordon and Daugherty 2003; Sisk and Mack 2018; van der Velden *et al.* 2021). In case of disclosure, oncologists' estimates are usually imprecise (e.g., "months to years") and framed optimistically (e.g., best-case scenarios), possibly to protect the patient (Chou *et al.* 2017; Derry *et al.* 2019a; Gordon and Daugherty 2003; Sisk and Mack 2018; van der Velden *et al.* 2021). The influence of oncologists' current communicative behavior on affective outcomes is relatively unknown (van der Velden *et al.* 2020).

Although emotional reactions to bad news are natural (Derry *et al.* 2019a), more insight into how oncologists' communicative behaviors affect patients is necessary to facilitate prognostic disclosure without doing unnecessary harm. This is important, as prognostic disclosure concerns sensitive and potentially confrontational information for patients in a vulnerable position, and negative emotions may inhibit patients' processing of information (Derry *et al.* 2019b). Indeed, researchers often appeal for honest, empathetic prognostic communication that balances realism with hope (Campbell *et al.* 2010; Parker *et al.* 2007; van Vliet *et al.* 2013a). In an era of patient-centeredness, not only the instrumental but also the affective purposes of medical communication are considered essential, satisfying patients' cognitive needs for information and emotional needs for support (van Vliet *et al.* 2013c). Yet, evidence-based guidelines for such prognostic communication are lacking, as recommendations are frequently expert opinion-based (Gilligan *et al.* 2017).

Existing studies cannot indisputably confirm or deny that prognostic disclosure versus non-disclosure evokes emotional suffering; evidence for influences on patients' appreciation of the consultation is limited (van der Velden *et al.* 2020). Little attention has been paid to how prognostic disclosure might affect patients' way of coping with cancer, while such coping influences patients' emotional adjustment to the disease and their psychological well-being (Johansson *et al.* 2011; Nipp *et al.* 2016). Notably, current knowledge mostly relies on non-causal associations with prognostic disclosure, and the impact of communicating unpredictability of prognosis seems unstudied altogether (van der Velden *et al.* 2020). Apart from the *types* of disclosure, little is known about effects of the *content* of oncologists' messages, when they do disclose prognosis. Qualitative research suggests that both vague word-based information and precise numbers could harm patients (Westendorp *et al.* 2022). Two experiments, however, found that numerical prognostic estimates decrease patients' uncertainty and increase satisfaction, without inducing anxiety (Mori *et al.* 2019; van Vliet *et al.* 2013c). Also, explaining multiples of the estimated group-level median overall survival as best-case, most likely and worst-case survival scenarios are presumed to be less upsetting and more hopeful than disclosing a median overall survival estimate only (Kiely *et al.* 2013).

The shift toward patient-centered care advocates tailoring prognostic information to individuals, given that patients may differ in their information preferences, ability to understand prognostic estimates, and personalities (Hoesseini *et al.* 2020). Indeed, effects of different prognostic messages could vary by moderating individual background characteristics. Scant evidence suggests that higher-educated patients particularly appreciate receiving standard, best- and worst-case survival scenarios (Kiely *et al.* 2013). Personal characteristics like dispositional optimism or uncertainty tolerance may shape patients' affective responses to prognostic communication too, as may patients' preferences for striving for

life-extension and for knowing prognosis (van der Velden *et al.* 2022a; van Vliet *et al.* 2013c). To illustrate, researchers found that patients' approaches to deal with threatening situations influence the effects of numerical estimates on patients' feelings of anxiety, uncertainty, and satisfaction (van Vliet *et al.* 2013c). While moving away from a one-size-fits-all approach seems relevant, scientific knowledge on *how* to best communicate prognosis to *whom* remains scarce (Gilligan *et al.* 2017; van der Velden *et al.* 2020).

When manipulating communication in reality is considered unethical, video-vignette studies offer a validated design that has been shown appropriate to demonstrate the effects of communication strategies on affective outcomes, including emotions (Mori *et al.* 2019; Nishioka *et al.* 2019; van Osch *et al.* 2014, 2017; van Vliet *et al.* 2013c; Visser *et al.* 2019). Video-vignettes are short visual depictions of scripted events (e.g., physician-patient conversations), rated by analogue patients (APs). APs are defined as patients, former patients, or healthy people who are instructed to imagine themselves in the shoes of the patient in the video (Hillen *et al.* 2013b, 2012; van Vliet *et al.* 2013b). Cancer-naïve APs can serve as valid proxies for patients from the general population (van Vliet *et al.* 2012), which prevents confounding influences of actual patients' prior personal experiences and burdening them with reliving a bad news conversation. Hence, we examined the independent, causal effects of various prognostic communication strategies in the advanced cancer setting on emotions (positive affect, negative affect, and state anxiety), coping (helplessness/hopelessness and fighting spirit), and appreciation of the consultation (trust in the oncologist; uncertainty, satisfaction, and desirability regarding provided prognostic information) among APs, *i.e.*:

- (1) The *type* of disclosure: prognostic disclosure vs. communication of unpredictability vs. non-disclosure.
- (2) The *content* of disclosure: *framing* (standard vs. standard and best-case vs. standard, best- and worst-case survival scenarios) and *precision* (numerical vs. word-based estimates) of prognosis.
- (3) The influence of individual characteristics (*i.e.*, background, personal, and patient preferences) on effects of the *type* and *content* of disclosure (see Table A.1, Appendix, for hypotheses).

These scientific efforts could provide preliminary data to inform evidence-based guidelines to navigate oncologists in the challenging task of prognostic communication.

## Methods

### Study design

The current paper reports on 1 of 2 sets of planned analyses on data collected in a randomized controlled experimental study, using video-vignettes of a simulated consultation (van der Velden *et al.* 2024). In our video-vignettes, professional actors portrayed a male patient with metastatic esophageal cancer, his daughter and a female oncologist, having a scripted discussion about treatment options with and without chemotherapy. Communication was standardized, yet we systematically manipulated the *type* and *content* of disclosure of prognosis.

We adopted a parallel group design to examine the *type* of disclosure: prognostic disclosure vs. communication of unpredictability vs. non-disclosure. We further manipulated the *content*

**Table 1.** Overview of conditions, including number of participants (van der Velden et al. 2024)

Manipulation: type of disclosure of prognosis		<i>n</i>	Duration of total video and variable content (minutes-seconds) <sup>a</sup>		
Parallel group design	Condition 1: Non-disclosure	208	7.23		
	Condition 2: Communication of unpredictability	210	7.54 (.31)		
Conditions 3–8: Prognostic disclosure	<b>Manipulation: content of disclosure of prognosis</b>				
	Balanced factorial 3 × 2 design		<b>Factor: framing of prognosis</b> (3 levels)	<b>Factor: precision of prognosis</b> (2 levels)	
		Condition 3	Standard scenarios	Numerical estimates	103
		Condition 4	Standard + best-case scenarios	Numerical estimates	103
		Condition 5	Standard + best-case + worst-case scenarios	Numerical estimates	100
		Condition 6	Standard scenarios	Word-based estimates	104
		Condition 7	Standard + best-case scenarios	Word-based estimates	103
		Condition 8	Standard + best-case + worst-case scenarios	Word-based estimates	105

*n* = sample size.

<sup>a</sup>The minutes and seconds between parentheses present the time of the video that includes variable content (i.e., manipulated prognostic communication).

of disclosure conditions in a balanced factorial design, combining 2 factors: *framing* with 3 levels (standard vs. standard and best-case vs. standard, best- and worst-case survival scenarios) and *precision* with 2 levels (numerical vs. word-based estimates). This led to a total of 8 conditions, as shown in Table 1. Figure A.1 (Appendix) presents a visual overview.

Numerical estimates were derived from published clinical trial data (Cunningham et al. 2008; Ter Veer et al. 2016). We used the group-level median overall survival for metastatic esophageal cancer with and without chemotherapy as standard scenarios and 90th and 10th percentiles of these curves as best- and worst-case scenarios, respectively (see Table A.2, Appendix, for the script including these manipulations).

### Participants

Individuals were eligible for participation if they were  $\geq 18$  years and had Dutch proficiency, internet access, no audiovisual impairments, and no self-reported (previous) cancer diagnosis.

### Study procedures

The institutional medical ethics review board provided exemption from formal approval (W20\_347 # 20.385app). Recruitment of APs and data collection were handled by an independent research agency (Flycatcher Internet Research). The agency stratified eligible APs in their panel by sex, age (5-year interval groups), and education prior to recruitment, to include a sample representative of our target population, i.e., Dutch patients with stage-IV cancer. Data from Netherlands Cancer Registry and Statistics Netherlands were used for stratification.

The research agency sent baseline surveys (T0) to the stratified sample in their private, secure, digital platform. APs who provided informed consent and completed T0 received an outcomes survey (T1) after  $\leq 3$  weeks. APs responded to items immediately before (T1, pre-video) and after (T1, post-video) watching 1 randomly assigned video. APs were allocated to 1 condition upon starting the video, using blocked randomization with a predetermined fixed size and sequence, i.e., 1122345678. An allocation ratio of 2:1 was used for conditions 1 and 2 (non-disclosure and communication of unpredictability) to enable detection of effects of the *type* of disclosure. APs were compensated within a digital reward system. The research agency executed data validity checks after finishing data collection.

### Development and piloting

Table A.3 (Appendix) describes development and piloting processes in detail, including example items and Cronbach's alphas.

### Study materials

Video-vignettes were developed by medical communication researchers and oncologists, consistent with published recommendations (Hillen et al. 2013b; van Vliet et al. 2013b). The script (see Table A.2, Appendix) was based on previously audio-recorded oncological consultations, past video-vignette studies, and existing communication guidelines (Clayton et al. 2007; Henselmans et al. 2017; Mori et al. 2019; van der Velden et al. 2021; van Vliet et al. 2013c).

Video-vignettes included an introductory waiting room scene, in which a voice-over conveys the backstory of the patient and

instructs APs to imagine themselves being in the depicted situation. In a subsequent consultation-scene, the oncologist discusses treatment options. Except the non-disclosure condition, all conditions included supplementary content, comprising either communication of unpredictability or prognostic disclosure for treatment with and without chemotherapy. We created a script with standard survival scenarios, which could be supplemented with best-case scenarios only or also with worst-case scenarios (*framing*), and developed versions with either numerical or word-based estimates (*precision*).

We evaluated manipulation success, credibility, medical content, and overall quality of the script with video-vignette experts and oncologists and organized rehearsals with professional actors. Next, we recorded and piloted 8 test videos. We evaluated manipulation success, credibility, comprehensibility, emotional load, and overall quality of test videos, using semi-structured interviews ( $n = 13$  cancer patients, 2 videos each) and online surveys ( $n = 120$  cancer-naïve individuals, 1 video each). After agreeing on an adjusted, final script, we professionally recorded 8 new videos and conducted semi-structured interviews with 5 more cancer patients for a “go or no-go” decision. Average video duration was 9 minutes and 3 seconds (range: 7.23–10.20 minutes/seconds).

### Surveys

Surveys were piloted among 8 cancer-naïve individuals ( $n = 2/8$  providing “think-aloud” feedback (Eccles and Aarsal 2017);  $n = 6/8$  providing feedback retrospectively) and pre-tested by the research agency.

### Study procedures

A subgroup of eligible APs took part in a *test phase* (April 2021;  $n = 41/79$  completed participation) to enable evaluation of study procedures ( $n = 14/41$  telephonic interviews). Hereupon, we slightly adjusted some survey instructions. As our study proved feasible, we proceeded with the *study phase* (May 2021;  $n = 1003/1828$  completed participation). Figure 1 presents an overview of our study.

### Measurements

Table A.4 (Appendix) describes all measures in detail.

### Individual characteristics

We measured the following background characteristics (T0): sex, age, religion, education, knowledge of cancer, and experience with oncological consultations for a loved one. Regarding personal characteristics, we assessed trait optimism (Life Orientation Test-Revised (Scheier et al. 1994)) and uncertainty tolerance (Tolerance for Ambiguity Scale (Geller et al. 1993)). After instructing APs to imagine being diagnosed with cancer, we measured patient preferences: prognostic information preference (“are you a person who wants to know your life expectancy? Yes/No” (Hagerty et al. 2004)) and attitude toward striving for length of life (Quality Quantity Questionnaire subscale (Stiggelbout et al. 1996)).

### Validity checks

To examine if manipulations were perceived as intended (T1, post-video), we assessed APs’ perceptions of the *type* and *content* of disclosure of prognosis. We also measured APs’ engagement with videos (shortened Video Engagement Scale [VES] (Lehmann et al. 2023) and perceived video realism (Fruijtjer et al. 2022; Hillen et al. 2013b; Lehmann et al. 2023; Medendorp et al. 2017, 2021a;

Visser et al. 2017, 2022, 2016a, 2016b, 2018). The shortened VES involves a screener item (i.e., “I was fully concentrated on the video while watching”; 1–7, “totally disagree” to “totally agree”); APs scoring  $\leq 2$  are excluded from analyses (Lehmann et al. 2023).

### Outcomes

Primary outcomes included APs’ *emotional reaction* (T1, pre-video; T1, post-video): differences in pre- and post-video scores on positive affect, negative affect (Positive and Negative Affect Scale (Watson et al. 1988a)), and state anxiety (Spielberger State-Trait Anxiety Inventory-Short Form State subscale (Marteau and Bekker 1992)). APs reported how they felt at that very moment (i.e., “now”).

For our secondary outcomes, we instructed APs to imagine being the depicted patient once again. We assessed APs’ *coping with cancer* (T1, post-video): helplessness/hopelessness (Mental Adjustment to Cancer subscale (Watson et al. 1988b)) and fighting spirit (Mini Mental Adjustment to Cancer subscale (Watson et al. 1994)). We also assessed APs’ *appreciation of the consultation* (T1, post-video): trust in the oncologist (Trust in Oncologist Scale-Short Form for APs (Hillen et al. 2013a, 2014, 2012, 2017)) and uncertainty (“the explanations this oncologist gives about my life expectancy seem hazy to me”; 1–5, “strongly disagree” to “strongly agree”), satisfaction (“how satisfied are you with the information you received from this oncologist about the general life expectancy of people with metastatic esophageal cancer?”; 0–10, “not at all satisfied” to “very satisfied”) and desirability (“would you like to receive information about life expectancy as was provided by the oncologist in the video?”; 1–5, “I certainly would not” to “I certainly would”) regarding provided prognostic information. Items on uncertainty, satisfaction, and desirability regarding provided prognostic information (adjusted from Mishel Uncertainty in Illness Scale (Mishel 1981; Mishel and Clayton 2008) and Patient Satisfaction Questionnaire (Aalfs et al. 2007)) were omitted from the non-disclosure condition.

### Statistical analysis

#### Power


Each of 6 prognostic disclosure conditions (videos 3–8; balanced factorial design) required 103 APs to detect main and moderating effects, based on a-priori G\*Power analyses ( $f = .125$ , small-medium size; Analysis of variance (ANOVA); power: 80%;  $\alpha = .05$ ;  $df = 2$ ). The non-disclosure and communication of unpredictability conditions (videos 1–2; parallel group design) each required 206 APs, resulting in a total sample of  $N = 1030$ .

#### Preparatory analyses

We checked scores for the screener item of the shortened VES. Manipulation success was examined by comparing proportions of APs who *believed* having watched a video with a certain manipulation among APs who *truly* watched a video with and without this manipulation ( $\chi^2$  tests). We tested differences between all 8 conditions for APs’ age, sex, and education ( $\chi^2$  tests, 1-way ANOVAs).

#### Primary analyses

To examine the independent effects of the *type* (prognostic disclosure vs. communication of unpredictability vs. non-disclosure) and *content* (*framing* and *precision*) of disclosure, we used separate multivariate analyses of variance (MANOVAs). For simultaneous significance testing of multiple outcomes, we clustered outcomes



**Figure 1.** Overview of phases and procedures of this study (van der Velden et al. 2024).

$n$  = sample size.<sup>a</sup>Based on existing audio-recorded oncological consultations, video-vignette studies, and prognostic communication guidelines.<sup>b</sup>Interviews were conducted among cancer patients (each watching 2 videos) and surveys among cancer-naive individuals (each watching 1 video).<sup>c</sup>Based on the content of the video-vignettes. Think-aloud and retrospective feedback procedures were conducted among cancer naive-individuals.<sup>d</sup> $n = 29/79$  (T0) and  $n = 9/50$  (T1) APs in the *test phase* and, respectively,  $n = 572/1828$  (T0) and  $n = 253/1256$  (T1) APs in the *study phase* were non-responders, drop-outs, or non-serious responders.<sup>e</sup> $n = 8/1044$  APs were excluded from the analyses (scores  $< 2$  on the Video Engagement Scale's screener item, i.e., "I was fully concentrated on the video while watching," "1: totally disagree" to "7: totally agree").

that were theoretically coherent and statistically correlated ( $r < .9$ ) (Tabachnick et al. 2007). This led to 3 clusters: (1) *emotional reaction* ( $\Delta$ positive affect,  $\Delta$ negative affect, and  $\Delta$ state anxiety, i.e., post-video scores minus pre-video scores;  $r = -.5$  to  $.7$ ); (2) *coping with cancer* (helplessness/hopelessness and fighting spirit;  $r = -.5$ ); and (3) *appreciation of the consultation* (trust in the oncologist and uncertainty, satisfaction, and desirability regarding provided prognostic information;  $r = -.5$  to  $.5$ ).

To investigate moderators, we added interaction terms of the *type of disclosure*, *framing*, and *precision* with APs' individual characteristics. To prevent multiple testing, we chose unique interaction terms for each cluster of outcomes based on predefined hypotheses (see Table A.1, Appendix). We conducted mean centering for continuous moderators.

To determine significance of MANOVAs' omnibus tests ( $\alpha = .05$ ), we evaluated Hotelling's Trace test-statistic, given its alleged robustness for violation of multivariate normality and homogeneity of covariance matrices (Field 2005). If significant, we evaluated MANOVAs' univariate follow-up tests, conducted post hoc analyses where necessary and reported statistics for models including moderators and interactions. We performed sensitivity analyses with Kruskal-Wallis and Mann-Whitney  $U$  tests for outcomes with distributions deviating from normality. Analyses were conducted with IBM-SPSS-26.

## Results

Of 1044 APs with complete data (response rate 55%; Figure A.2), we excluded 8 individuals with the shortened VES' screener item

(Lehmann et al. 2023). Of 1036 APs in the analyses, 60% was male. Age ranged between 21 and 89 years ( $M_{\text{age}} = 66$ ). The majority of APs (63%) reported their cancer knowledge to be average. APs' engagement with and perceived realism of videos were satisfactory ( $M_{\text{Video engagement}} = 5.3$ ,  $M_{\text{Video realism}} = 5.9$ ). Table 2 describes APs' background characteristics.

Manipulations were perceived as intended, although *framing* was less clearly perceived by APs than other manipulations, as shown in Table A.5. Table 3 describes the potential moderating characteristics (see Table A.1, Appendix, for hypotheses), validity checks and outcomes, stratified by manipulation. There were no significant differences in APs' sex, age, or education between conditions.

### The type of disclosure

Tables A.6–A.9 (Appendix) demonstrate all results. We found no significant overall effect of the type of disclosure on *emotional reaction*, nor significant moderating effects of APs' *prognostic information preference* or *uncertainty tolerance*.

There was no significant overall effect of the type of disclosure on *coping with cancer*, nor significant moderating effects of APs' *trait optimism* or *attitude toward striving for length of life*.

We found a significant overall effect of the type of disclosure on *appreciation of the consultation* ( $F(3, 807) = 3.932$ ,  $p_{\text{omnibus}} = .008$ , Hotelling's Trace =  $.015$ ). Univariate results showed that APs reported more *satisfaction* ( $F(1, 809) = 11.303$ ,  $p < .001$ ) with provided prognostic information after prognostic disclosure, compared to communication of unpredictability.

**Table 2.** Background characteristics of the total sample<sup>a</sup>

APs' characteristics	Total sample <i>n</i> = 1036
Sex (male), % ( <i>n</i> )	60.1 (623)
Age (years), mean ± SD	65.77 ± 10.85
Education, % ( <i>n</i> ) <sup>b</sup>	
Low	29.2 (303)
Medium	42.0 (435)
High	28.8 (298)
Religion, % ( <i>n</i> )	
Christianity	44.3 (459)
Other	6.2 (64)
None	49.5 (513)
Knowledge of cancer, % ( <i>n</i> )	
None or very little	4.3 (45)
Little	24.9 (258)
Average	63.1 (654)
Much	7.1 (74)
Very much	.5 (5)
Experience with oncological consultations for a loved one (yes), % ( <i>n</i> )	36.4 (377)
Trait optimism (LOT-R, 0–24), mean ± SD	14.96 ± 3.48
Uncertainty tolerance (TFA, 7–42), mean ± SD	24.53 ± 5.02
Prognostic information preference (yes), % ( <i>n</i> )	90.7 (940)
Attitude toward striving for length of life (QQQ, 4–20), mean ± SD	11.76 ± 3.31
Video engagement (shortened VES, 1–7), mean ± SD	5.31 ± 1.15
Perceived video realism (1–7), mean ± SD	5.90 ± 1.04

AP = analogue patient; *n* = sample size; SD = standard deviation; LOT-R = Life Orientation Test-Revised; TFA = Tolerance for Ambiguity Scale; QQQ = Quality Quantity Questionnaire; VES = Video Engagement Scale.

<sup>a</sup>The content of this table partly overlaps results of a different analysis (reported elsewhere (van der Velden et al. 2024)) on the dataset used for the current paper.

<sup>b</sup>Low vocational education; medium level vocational education; high vocational or academic education.

Confirming our hypotheses, we found a significant moderating effect of *prognostic information preference* ( $F(3, 807) = 3.451$ ,  $p_{\text{omnibus}} = .016$ , Hotelling's Trace = .013). APs who preferred to know prognosis reported less *uncertainty* and more *desirability* regarding provided prognostic information after prognostic disclosure than communication of unpredictability, whereas APs who preferred not knowing reported less *uncertainty* and more *desirability* after communication of unpredictability instead (interactions:  $F_{\text{uncertainty}}(1, 809) = 7.291$ ,  $p = .007$ ;  $F_{\text{desirability}}(1, 809) = 6.268$ ,  $p = .012$ ).

Furthermore, we found a significant moderating effect of *uncertainty tolerance* ( $F(3, 807) = 2.794$ ,  $p_{\text{omnibus}} = .039$ , Hotelling's Trace = .010). Opposing our hypotheses, APs with higher uncertainty tolerance considered information more *desirable* after prognostic disclosure than communication of unpredictability, whereas APs with lower uncertainty tolerance considered communication of unpredictability more *desirable* instead

(interaction:  $F(1, 809) = 7.811$ ,  $p = .005$ ). Also contradicting our hypotheses, APs with higher uncertainty tolerance reported more *trust* in the oncologist after prognostic disclosure than non-disclosure, while APs with lower uncertainty tolerance reported more *trust* after non-disclosure instead (interaction:  $F(2, 1027) = 3.350$ ,  $p = .035$ ).

### The content of disclosure

There were no significant overall effects of framing or the precision of prognosis on *emotional reaction*. We found a significant interaction between APs' *uncertainty tolerance* and precision ( $F(3, 608) = 3.233$ ,  $p_{\text{omnibus}} = .022$ , Hotelling's Trace = .016). Opposing our hypotheses, yet in line with aforementioned moderating effects, APs with higher uncertainty tolerance reported more *positive affect*, less *negative affect*, and less *state anxiety* after numerical rather than word-based estimates. APs with lower uncertainty tolerance reported more *positive affect*, less *negative affect*, and less *state anxiety* after word-based estimates instead (interactions:  $F_{\text{positive affect}}(1, 610) = 5.983$ ,  $p = .015$ ;  $F_{\text{negative affect}}(1, 610) = 7.867$ ,  $p = .005$ ;  $F_{\text{state anxiety}}(1, 610) = 5.870$ ,  $p = .016$ ).

We found no significant overall effects of framing or the precision of prognosis on *coping with cancer*, nor significant moderating effects of APs' *trait optimism* or *attitude toward striving for length of life*.

There was no significant overall effect of framing on *appreciation of the consultation*. However, we found a significant overall effect of precision ( $F(4, 593) = 2.978$ ,  $p_{\text{omnibus}} = .019$ , Hotelling's Trace = .020). Univariate results showed that APs reported less *uncertainty* ( $F(1, 596) = 4.139$ ,  $p = .042$ ), more *satisfaction* ( $F(1, 596) = 7.791$ ,  $p = .005$ ), and more *desirability* ( $F(1, 596) = 5.792$ ,  $p = .016$ ) regarding provided prognostic information after numerical instead of word-based estimates. APs' *education* or *uncertainty tolerance* did not moderate effects.

All aforementioned findings were confirmed in non-parametric sensitivity analyses.

### Discussion

Prognostic information is important for patients with advanced cancer to accustom to the last phase of life (Butow et al. 2020; van der Velden et al. 2020). Disclosing prognosis, however, is challenging for oncologists and confrontational for patients (Franssen et al. 2009; Liu et al. 2014). This study examined the effects of oncologists' prognostic messages on affective outcomes, hereby advancing scientific guidance on treading the fine line between disclosing poor prospects and preserving patients' emotional well-being. Overall, disclosure of (precise) prognostic estimates did not increase feelings of anxiety, negativity, or hopelessness nor decrease feelings of positivity or fighting spirit among APs. On average, individuals were more satisfied with prognostic disclosure than communication of unpredictability and generally considered numbers more satisfactory, desirable, and certain than word-based estimates. However, importantly, some of these effects depended on individuals' prognostic information preference and uncertainty tolerance. Communication of best- and/or worst-case survival scenarios lacked effects on all outcomes.

Our finding that individuals with a preference for not knowing prognosis (comprising 9% of the sample) were less appreciative of prognostic disclosure makes sense, as they might feel that their needs were not met. Inherent to our design, the oncologist in the video disclosed prognosis irrespective of the

**Table 3.** Hypothesized moderators, validity checks, and outcomes (stratified by manipulation)<sup>a</sup>

	Non-disclosure (video 1) <i>n</i> = 208	Communication of unpredictability (video 2) <i>n</i> = 210	Prognostic disclosure (videos 3–8) <i>n</i> = 618	Prognostic disclosure with standard and best-case scenarios (videos 3 and 6) <i>n</i> = 207	Prognostic disclosure with standard and best-case scenarios (videos 4 and 7) <i>n</i> = 206	Prognostic disclosure with standard, best- and worst-case scenarios (videos 5 and 8) <i>n</i> = 205	Prognostic disclosure with numerical estimates (videos 3–5) <i>n</i> = 306	Prognostic disclosure with word-based estimates (videos 6–8) <i>n</i> = 312
<b>Hypothesized moderators</b>								
Education, % ( <i>n</i> ) <sup>b</sup>								
Low	26.4 (55)	28.1 (59)	30.6 (189)	31.4 (65)	33.5 (69)	26.8 (55)	27.5 (84)	33.7 (105)
Medium	43.3 (90)	45.7 (96)	40.3 (249)	41.1 (85)	39.3 (81)	40.5 (83)	40.5 (124)	40.1 (125)
High	30.3 (63)	26.2 (55)	29.1 (180)	27.5 (57)	27.2 (56)	32.7 (67)	32.0 (98)	26.3 (82)
Trait optimism (LOT-R, 0–24), mean ± SD	14.78 ± 3.40	14.47 ± 3.27	15.18 ± 3.55	14.92 ± 3.55	15.42 ± 3.56	15.20 ± 3.56	15.27 ± 3.61	15.09 ± 3.50
Uncertainty tolerance (TFA, 7–42), mean ± SD	24.34 ± 5.20	24.49 ± 4.86	24.61 ± 5.01	24.68 ± 4.86	24.61 ± 4.92	24.56 ± 5.26	24.85 ± 5.25	24.38 ± 4.76
Prognostic information preference (yes), % ( <i>n</i> )	92.8 (193)	93.3 (196)	89.2 (551)	86.5 (179)	90.8 (187)	90.2 (185)	89.2 (273)	89.1 (278)
Attitude toward striving for length of life (QQQ subscale, 4–20), mean ± SD	11.60 ± 3.29	12.23 ± 3.22	11.65 ± 3.34	11.61 ± 3.33	11.90 ± 3.17	11.43 ± 3.52	11.53 ± 3.30	11.76 ± 3.38
<b>Validity checks</b>								
Video engagement (shortened VES, 1–7), mean ± SD	5.23 ± 1.18	5.29 ± 1.16	5.35 ± 1.14	5.27 ± 1.15	5.34 ± 1.18	5.45 ± 1.09	5.35 ± 1.15	5.35 ± 1.13
Perceived video realism (1–7), mean ± SD	5.86 ± .99	5.83 ± 1.17	5.94 ± 1.00	5.81 ± 1.06	5.92 ± .97	6.09 ± .95	5.95 ± .99	5.92 ± 1.01

(Continued)

Table 3. (Continued.)

	Non-disclosure (video 1) <i>n</i> = 208	Communication of unpredictability (video 2) <i>n</i> = 210	Prognostic disclosure (videos 3–8) <i>n</i> = 618	Prognostic disclosure with standard scenarios (videos 3 and 6) <i>n</i> = 207	Prognostic disclosure with standard and best-case scenarios (videos 4 and 7) <i>n</i> = 206	Prognostic disclosure with standard, best- and worst-case scenarios (videos 5 and 8) <i>n</i> = 205	Prognostic disclosure with numerical estimates (videos 3–5) <i>n</i> = 306	Prognostic disclosure with word-based estimates (videos 6–8) <i>n</i> = 312
<b>Outcomes</b>								
ΔPositive affect (PANAS subscale, 0–40), mean ± SD <sup>c</sup>	-5.13 ± 5.75	-5.41 ± 6.20	-5.90 ± 6.17	-5.88 ± 6.31	-5.80 ± 6.47	-6.02 ± 5.73	-6.16 ± 5.86	-5.64 ± 6.46
ΔNegative affect (PANAS subscale, 0–40), mean ± SD <sup>c</sup>	4.34 ± 5.97	4.10 ± 6.40	5.18 ± 6.42	5.03 ± 6.30	4.96 ± 6.49	5.57 ± 6.49	4.97 ± 6.52	5.39 ± 6.33
ΔState anxiety (STAI-state-SF, 0–18), mean ± SD <sup>c</sup>	2.65 ± 4.07	2.59 ± 3.91	3.02 ± 4.13	2.76 ± 4.10	3.16 ± 4.30	3.15 ± 3.99	2.92 ± 4.11	3.12 ± 4.15
Helplessness/hopelessness (MAC subscale, 6–24), mean ± SD	14.95 ± 2.86	14.91 ± 2.69	14.88 ± 3.05	14.89 ± 3.05	14.66 ± 3.13	15.09 ± 2.98	14.88 ± 2.91	14.88 ± 3.19
Fighting spirit (mini- MAC subscale, 4–16), mean ± SD	9.99 ± 2.40	10.13 ± 2.20	10.23 ± 2.35	10.01 ± 2.28	10.45 ± 2.36	10.22 ± 2.40	10.20 ± 2.34	10.26 ± 2.37
Trust in oncologist (TIOS-SF, 1–5), mean ± SD	4.06 ± .51	4.11 ± .54	4.19 ± .51	4.13 ± .54	4.19 ± .48	4.24 ± .49	4.19 ± .51	4.18 ± .51

(Continued)



**Table 3.** (Continued.)

	Non-disclosure (video 1) n = 208	Communication of unpredictability (video 2) n = 210	Prognostic disclosure (videos 3–8) n = 618	Prognostic disclosure with standard scenarios (videos 3 and 6) n = 207	Prognostic disclosure with standard and best-case scenarios (videos 4 and 7) n = 206	Prognostic disclosure with standard, best- and worst-case scenarios (videos 5 and 8) n = 205	Prognostic disclosure with numerical estimates (videos 3–5) n = 306	Prognostic disclosure with word-based estimates (videos 6–8) n = 312
Uncertainty regarding prognostic information (MUIS selected items, 1–5), mean ± SD <sup>d,e</sup>	x 2.50 ± .78 <sup>f</sup>	2.11 ± .76 <sup>g</sup>	2.17 ± .77 <sup>h</sup>	2.10 ± .72 <sup>h</sup>	2.07 ± .78	2.04 ± .77 <sup>h</sup>	2.18 ± .74 <sup>h</sup>	
Satisfaction regarding prognostic information (PSQ, selected item, 0–10), mean ± SD <sup>d,e</sup>	x 6.27 ± 2.08 <sup>f</sup>	7.48 ± 1.61 <sup>g</sup>	7.30 ± 1.61 <sup>h</sup>	7.54 ± 1.54 <sup>h</sup>	7.61 ± 1.65	7.64 ± 1.54 <sup>h</sup>	7.32 ± 1.66 <sup>h</sup>	
Desirability regarding prognostic information (1–5), mean ± SD <sup>d</sup>	x 3.58 ± .97 <sup>f</sup>	4.11 ± .85 <sup>g</sup>	3.97 ± .88 <sup>h</sup>	4.21 ± .80 <sup>h</sup>	4.17 ± .83	4.20 ± .84 <sup>h</sup>	4.03 ± .84 <sup>h</sup>	

n = sample size; SD = standard deviation; LOT-R = Life Orientation Test-Revised; TFA = Tolerance for Ambiguity Scale; QQQ = Quality Quantity Questionnaire; VES = Video Engagement Scale; PANAS = Positive and Negative Affect Scale; STAI-state-SF = State-subscale of the Spielberger State-Trait Anxiety Inventory-Short Form; MAC = Mental Adjustment to Cancer; TIOS-SF = Trust in Oncologist Scale-Short Form; MUIS = Mishel Uncertainty in Illness Scale; PSQ = Patient Satisfaction Questionnaire.

<sup>a</sup>The content of this table partly overlaps results of a different analysis (reported elsewhere (van der Velden et al. 2024)) on the dataset used for the current paper.

<sup>b</sup>Low vocational education; medium level vocational education; high vocational or academic education.

<sup>c</sup>Difference scores were calculated by subtracting pre-video scores from post-video scores on positive affect (PANAS subscale), negative affect (PANAS subscale), and state anxiety (STAI-STATE-SF). Negative difference scores indicate reduced positive affect, negative affect, and state anxiety, respectively. Positive difference scores indicate increased positive affect, negative affect, and state anxiety, respectively.

<sup>d</sup>Items were not presented to APs in the condition with non-disclosure (video 1).

<sup>e</sup>Original (selected) items of the MUIS and PSQ were adjusted for this study's purpose.

<sup>f</sup>7 missings (n = 203/210 for communication of unpredictability).

<sup>g</sup>6 missings (n = 612/618 for prognostic disclosure).

<sup>h</sup>3 missings (n = 204/207 for disclosure with standard scenarios only; n = 203/206 for disclosure with standard and best-case scenarios; n = 303/306 for disclosure with numerical estimates; n = 309/312 for disclosure with word-based estimates). Missings occurred if APs indicated having watched a video without communication about prognosis.

(depicted or imagined) patient's information preference. In clinical practice, overwhelming patients with more information than desired must be prevented at all times to avoid undue distress (Lehmann *et al.* 2020). Our finding that disclosing (precise) prognostic information generally has no impact on individuals' emotions or way of coping within an experimental setting, and that numbers yield more appreciation than words, resonates with van Vliet *et al.* (2013c), and with Mori *et al.* (2019). These video-vignette studies similarly found that numerical estimates do not evoke anxiety, yet do increase patients' certainty and satisfaction (Mori *et al.* 2019; van Vliet *et al.* 2013c). Our interaction analyses revealed that prognostic disclosure leads to less desirability of information and trust in the oncologist, and numbers lead to more anxious, more negative, and less positive feelings in individuals who poorly tolerate uncertainty. Although these results contradict our hypotheses, the moderating effects of uncertainty tolerance show similar directions and, in fact, apparently corroborate previous research (van Vliet *et al.* 2013c). Van Vliet *et al.* (2013c) found that numerical estimates increase anxiety and uncertainty and decrease satisfaction among "high monitors," *i.e.*, patients who, in the face of threat, seek ways to reduce uncertainty (van Vliet *et al.* 2013c). We may speculate that individuals who dislike uncertainty do not benefit from prognostic estimates, because predictions about life expectancy are fundamentally uncertain (Han *et al.* 2017). More precise and/or extensive information does not necessarily decrease such uncertainty. Inspecting our manipulations, we may pose that the multiplicity of numerical estimates (*e.g.*, survival with and without chemotherapy, survival gain, best-cases, and worst-cases) perhaps introduced complexity, which only added uncertainty (Han *et al.* 2017). Altogether, prognostic estimates possibly prompt more questions, which is unsettling to individuals with low uncertainty tolerance (Han *et al.* 2009; Rains and Tukachinsky 2015; Street *et al.* 2012). Opposed to individuals with low uncertainty tolerance, individuals with a preference for knowing prognosis (comprising 91% of the sample) particularly appreciated prognostic disclosure, which implies that those who poorly tolerate uncertainty are not necessarily the same individuals as those who seek prognostic information. As researchers theorize that low uncertainty tolerance fuels the appraisal of threat, exacerbating distress, it could be that those individuals benefit from more affective communication (*e.g.*, addressing emotions and reassuring continued care), alleviating distress, rather than more precise information (Chasiotis *et al.* 2021; Reis-Dennis *et al.* 2021; Taha *et al.* 2014).

Apart from aforementioned moderating effects, there was insufficient evidence for many predefined hypotheses about moderators. For example, opposing our expectations, individuals' educational level did not influence responses to numerical estimates or different survival scenarios, which disaffirms the conception that highly educated patients specifically value more detailed prognostic information (Franssen *et al.* 2009; Kiely *et al.* 2013; Loiselle 2019; Mayer *et al.* 2007).

Our research has several limitations. First, our methodological choices engender limited ecological validity. For ethical reasons, we included disease-naïve individuals, who are valid proxies for actual patients according to a previous meta-analysis (van Vliet *et al.* 2012). While scores for perceived video realism and video engagement were satisfactory in the current study, watching a video-vignette clearly does not equal being in an actual consultation and cancer-naïve individuals do not experience the physical, emotional, and existential burdens of advanced cancer. Hence, it is plausible that cancer-naïve individuals may react

less emotionally than patients, who need to reconcile their own existence with the inevitable state of nonexistence upon death. Clinical research previously observed mixed effects of prognostic disclosure on emotional well-being (Cripe *et al.* 2012; Enzinger *et al.* 2015; Fletcher *et al.* 2013; Nakajima *et al.* 2013; Shin *et al.* 2016), but those studies investigated delayed instead of immediate emotional responses and could not infer causality. Thus, although we recognize that extrapolation of our findings to clinical practice is speculative, we believe that manipulation of prognostic communication in a more ecologically valid way would not be ethically justified without at least some substantiation from experimental research with APs, especially since current evidence for this hard-to-study topic is at a preliminary stage. Future research can build on this video-vignette experiment through replication among individuals with a cancer diagnosis or a history of cancer to enhance real-world impact. Using more ecologically valid methods, virtual reality technologies allow systematic testing of prognostic communication elements within more realistic settings. Experiments may ultimately inform an intervention addressing prognostic communication skills for oncologists; its effectiveness could be tested in a clinical RCT. Researchers could audiotape oncologist-patient conversations to analyze whether the prognostic communication skills taught in the intervention are endorsed. A second limitation may be our manipulation of framing. Sixty percent of participants reported to recall additional survival scenarios on top of the standard scenarios, while viewing videos without such information (Table A.5). Perhaps, our standard survival scenarios essentially reflected a best-/worst-case format (*e.g.*, "months to a few years"). More research is necessary to substantiate evidence on different survival scenarios. Third, we did not establish validity of items that were tailored to this study, *i.e.*, appreciation of the consultation, nor did we determine thresholds for the clinical importance of statistical differences in our outcomes. Finally, given that prognostic communication is considered culturally sensitive, generalizability of our results may be restricted by the Dutch study setting. The Dutch culture places value on knowledge, autonomy, and straightforwardness. It is conceivable that patients from non-Western countries hold different attitudes toward prognostic communication (Surbone 2008). Unfortunately, our study lacks insight into participants' demographic information like race, ethnicity, or immigration status. Also regarding systematic disparities in health care, it is important to note that we used (biological) sex-disaggregated data, which do not consider gender identity. Strengths include our ability to draw causal conclusions about the independent effects of prognostic communication strategies, extensive pilot-testing procedures, and use of a large, stratified sample.

## Conclusion

Based on this experimental study, we tentatively conclude that the disclosure of prognosis does not necessarily cause more emotional harm than non-disclosure. Prognostic disclosure is particularly appreciated by patients who prefer to receive such information and those who tolerate uncertainty well. Generally, numerical estimates yield more appreciation than prognostic information in words. Yet, numerical estimates may evoke more negative emotions in patients with low uncertainty tolerance. When providing prognostic information in clinical consultations, it remains crucial that oncologists acknowledge and act on individual variation in patient preferences and personal characteristics.

### Practice implications

Responding to oncologists' discomfort with communicating prognosis, our results may potentially relieve worries about undermining hope or causing emotional harm. Despite the preliminary nature of findings, this study stipulates that oncologists' fears must not preclude offering prognostic information. In line with previous literature and our findings on moderating effects of information preference, we advocate firstly asking patients if they want prognostic information, as well as assessing their preferences for the content and timing of disclosure (Clayton et al. 2007; Gilligan et al. 2017; Lehmann et al. 2020; van der Velden et al. 2022b).

Our observation that prognostic disclosure versus communication of unpredictability and numerical versus word-based estimates generally promote satisfaction may stimulate provision of precise prognostic information. When navigating the delicate balance between realism and hope (Olsman et al. 2014), oncologists could allude to explaining different survival scenarios. Although the added value of this communication strategy remains unclear, it may be reassuring that disclosure of worst-case scenarios, in addition to best-case scenarios, is not necessarily upsetting. Importantly, our study does not imply that patients will not be moved by bad news. It is crucial for oncologists to allow space for emotions, listen and acknowledge patient concerns with empathy (Back and Arnold 2006; Westendorp et al. 2022).

Even more so, our results suggest sensitivity as to *which* prognostic communication strategies are beneficial for *whom*. Uncertainty tolerance appears to play an important role in how individuals react to prognostic information. It, therefore, seems valuable for oncologists to explore how patients generally deal with uncertainty, address the inherent uncertainty of prognostic estimates, and support patients in coping with prognostic uncertainty (Medendorp et al. 2021b). Oncologists could, for example, accentuate controllable elements of an uncertain situation by making concrete follow-up plans and explicating their continued involvement in patient's care (Medendorp et al. 2021b). Future research must further clarify the mechanisms underlying the association between prognostic communication and uncertainty tolerance. Additionally, while we found no moderating effects of education, trait optimism, and attitude toward striving for length of life, more studies should increase knowledge on tailoring prognostic communication.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S1478951524000403>.

**Acknowledgments.** The authors thank Onderwater Productions en Whaleshark Productions for developing the video-vignettes. The authors furthermore thank the individuals who took part in this study's pilot tests and data collection.

**Funding.** This work was supported by the Dutch Cancer Society (grant number 11547).

**Competing interests.** The authors declare no conflicts of interest.

### References

- Aalfs C, Oort F, De Haes J, et al. (2007) A comparison of counsellor and counselor satisfaction in reproductive genetic counseling. *Clinical Genetics* 72(2), 74–82. doi:10.1111/j.1399-0004.2007.00834.x
- Back AL and Arnold RM (2006) Discussing prognosis: "How much do you want to know?" talking to patients who do not want information or who

are ambivalent. *Journal of Clinical Oncology* 24(25), 4214–4217. doi:10.1200/JCO.2006.06.008

Butow PN, Clayton JM and Epstein RM (2020) Prognostic awareness in adult oncology and palliative care. *Journal of Clinical Oncology* 38(9), 877–884. doi:10.1200/JCO.18.02112

Campbell TC, Carey EC, Jackson VA, et al. (2010) Discussing prognosis: Balancing hope and realism. *The Cancer Journal* 16(5), 461–466. doi:10.1097/PPO.0b013e3181f30e07

Chasiotis A, Wedderhoff O, Rosman T, et al. (2021) The role of approach and avoidance motivation and emotion regulation in coping via health information seeking. *Current Psychology* 40, 5235–5244. doi:10.1007/s12144-019-00488-3

Chou WS, Hamel LM, Thai CL, et al. (2017) Discussing prognosis and treatment goals with patients with advanced cancer: A qualitative analysis of oncologists' language. *Health Expectations* 20(5), 1073–1080. doi:10.1111/hex.12549

Clayton JM, Hancock KM, Butow PN, et al. (2007) Clinical practice guidelines for communicating prognosis and end-of-life issues with adults in the advanced stages of a life-limiting illness, and their caregivers. *Medical Journal of Australia* 186(12 Suppl), S77–S105. doi:10.5694/j.1326-5377.2007.tb01100.x

Cripe LD, Rawl SM, Schmidt KK, et al. (2012) Discussions of life expectancy moderate relationships between prognosis and anxiety or depression in men with advanced cancer. *Journal of Palliative Medicine* 15(1), 99–105. doi:10.1089/jpm.2011.0249

Cunningham D, Starling N, Rao S, et al. (2008) Capecitabine and oxaliplatin for advanced esophagogastric cancer. *New England Journal of Medicine* 358(1), 36–46. doi:10.1056/NEJMoa073149

Daugherty CK and Hlubocky FJ (2008) What are terminally ill cancer patients told about their expected deaths? A study of cancer physicians' self-reports of prognosis disclosure. *Journal of Clinical Oncology* 26(36), 5988. doi:10.1200/JCO.2008.17.2221

Derry HM, Epstein AS, Lichtenthal WG, et al. (2019a) Emotions in the room: Common emotional reactions to discussions of poor prognosis and tools to address them. *Expert Review of Anticancer Therapy* 19(8), 689–696. doi:10.1080/14737140.2019.1651648

Derry HM, Reid MC and Prigerson HG (2019b) Advanced cancer patients' understanding of prognostic information: Applying insights from psychological research. *Cancer Medicine* 8(9), 4081–4088. doi:10.1002/cam4.2331

Eccles DW and Aarsal G (2017) The think aloud method: What is it and how do I use it? *Qualitative Research in Sport, Exercise and Health* 9(4), 514–531. doi:10.1080/2159676X.2017.1331501

Enzinger AC, Zhang B, Schrag D, et al. (2015) Outcomes of prognostic disclosure: Associations with prognostic understanding, distress, and relationship with physician among patients with advanced cancer. *Journal of Clinical Oncology* 33(32), 3809–3816. doi:10.1200/JCO.2015.61.9239

Field A (2005) *Discovering Statistics Using SPSS*. London: Sage publications Ltd.

Fletcher K, Prigerson HG, Paulk E, et al. (2013) Gender differences in the evolution of illness understanding among patients with advanced cancer. *The Journal of Supportive Oncology* 11(3), 126. doi:10.12788/j.suponc.0007

Franssen SJ, Lagarde SM, Van Werven JR, et al. (2009) Psychological factors and preferences for communicating prognosis in esophageal cancer patients. *Psychooncology* 18(11), 1199–1207. doi:10.1002/pon.1485

Fruijtjer AD, van der Schaar J, van Maurik IS, et al. (2022) Identifying best practices for disclosure of amyloid imaging results: A randomized controlled trial. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association* 19(1), 285–295.

Geller G, Tambor ES, Chase GA, et al. (1993) Measuring physicians' tolerance for ambiguity and its relationship to their reported practices regarding genetic testing. *Medical Care* 31(11), 989–1001. doi:10.1097/00005650-199311000-00002

Gilligan T, Coyle N, Frankel RM, et al. (2017) Patient-clinician communication: American Society of Clinical Oncology consensus guideline. *Journal of Clinical Oncology* 35(31), 3618–3632. doi:10.1200/JCO.2017.75.2311

Gordon EJ and Daugherty CK (2003) 'Hitting you over the head': Oncologists' disclosure of prognosis to advanced cancer patients. *Bioethics* 17(2), 142–168. doi:10.1111/1467-8519.00330

- Hagerty RG, Butow PN, Ellis PA, *et al.* (2004) Cancer patient preferences for communication of prognosis in the metastatic setting. *Journal of Clinical Oncology* 22(9), 1721–1730. doi:10.1200/JCO.2004.04.095pii
- Hancock K, Clayton JM, Parker SM, *et al.* (2007) Truth-telling in discussing prognosis in advanced life-limiting illnesses: A systematic review. *Palliative Medicine* 21(6), 507–517. doi:10.1177/0269216307080823
- Han PK, Klein WM, Lehman TC, *et al.* (2009) Laypersons' responses to the communication of uncertainty regarding cancer risk estimates. *Medical Decision Making* 29(3), 391–403. doi:10.1177/0272989X08327396
- Han PK, Umstead KL, Bernhardt BA, *et al.* (2017) A taxonomy of medical uncertainties in clinical genome sequencing. *Genetics in Medicine* 19(8), 918–925. doi:10.1038/gim.2016.212
- Henselmans I, Smets EMA, Han PKJ, *et al.* (2017) How long do I have? Observational study on communication about life expectancy with advanced cancer patients. *Patient Education and Counseling* 100(10), 1820–1827. doi:10.1016/j.pec.2017.05.012
- Hillen MA, Butow PN, Tattersall MH, *et al.* (2013a) Validation of the English version of the Trust in Oncologist Scale (TiOS). *Patient Education and Counseling* 91(1), 25–28. doi:10.1016/j.pec.2012.11.004
- Hillen MA, de Haes HCJM, Stalpers LJA, *et al.* (2014) How can communication by oncologists enhance patients' trust? An experimental study. *Annals of Oncology* 25(4), 896–901. doi:10.1093/annonc/mdu027
- Hillen MA, Koning CCE, Wilmink JW, *et al.* (2012) Assessing cancer patients' trust in their oncologist: Development and validation of the Trust in Oncologist Scale (TiOS). *Supportive Care in Cancer* 20(8), 1787–1795. doi:10.1007/s00520-011-1276-8
- Hillen MA, Postma R-M, Verdam MGE, *et al.* (2017) Development and validation of an abbreviated version of the Trust in Oncologist Scale—The Trust in Oncologist Scale—short form (TiOS-SF). *Supportive Care in Cancer* 25(3), 855–861. doi:10.1007/s00520-016-3473-y
- Hillen MA, van Vliet LM, de Haes HC, *et al.* (2013b) Developing and administering scripted video vignettes for experimental research of patient-provider communication. *Patient Education and Counseling* 91(3), 295–309. doi:10.1016/j.pec.2013.01.020
- Hoesseini A, Dronkers EA, Sewnaik A, *et al.* (2020) Head and neck cancer patients' preferences for individualized prognostic information: A focus group study. *BMC Cancer* 20, 1–14. doi:10.1186/s12885-020-6554-8
- Innes S and Payne S (2009) Advanced cancer patients' prognostic information preferences: A review. *Palliative Medicine* 23(1), 29–39. doi:10.1177/0269216308098799
- Johansson M, Rydén A and Finizia C (2011) Mental adjustment to cancer and its relation to anxiety, depression, HRQL and survival in patients with laryngeal cancer – a longitudinal study. *BMC Cancer* 11(1), 283 doi:10.1186/1471-2407-11-283
- Kiely BE, McCaughan G, Christodoulou S, *et al.* (2013) Using scenarios to explain life expectancy in advanced cancer: Attitudes of people with a cancer experience. *Supportive Care in Cancer* 21(2), 369–376. doi:10.1007/s00520-012-1526-4
- Lehmann V, Hillen MA, Verdam MG, *et al.* (2023) The Video Engagement Scale (VES): Measurement properties of the full and shortened VES across studies. *International Journal of Social Research Methodology* 26, 305–318. doi:10.1080/13645579.2022.2052697
- Lehmann V, Labrie NH, van Weert JC, *et al.* (2020) Tailoring the amount of treatment information to cancer patients' and survivors' preferences: Effects on patient-reported outcomes. *Patient Education and Counseling* 103(3), 514–520. doi:10.1016/j.pec.2019.09.024
- Liu PH, Landrum MB, Weeks JC, *et al.* (2014) Physicians' propensity to discuss prognosis is associated with patients' awareness of prognosis for metastatic cancers. *Journal of Palliative Medicine* 17(6), 673–682. doi:10.1089/jpm.2013.0460
- Loisel CG (2019) Cancer information-seeking preferences linked to distinct patient experiences and differential satisfaction with cancer care. *Patient Education and Counseling* 102(6), 1187–1193. doi:10.1016/j.pec.2019.01.009
- Marteau TM and Bekker H (1992) The development of a six-item short-form of the state scale of the Spielberger State-Trait Anxiety Inventory (STAI). *The British Journal of Clinical Psychology* 31(3), 301–306. doi:10.1111/j.2044-8260.1992.tb00997.x
- Mayer DK, Terrin NC, Kreps GL, *et al.* (2007) Cancer survivors information seeking behaviors: A comparison of survivors who do and do not seek information about cancer. *Patient Education and Counseling* 65(3), 342–350. doi:10.1016/j.pec.2006.08.015
- Medendorp NM, Hillen MA, Visser LN, *et al.* (2021a) A randomized experimental study to test the effects of discussing uncertainty during cancer genetic counseling: Different strategies, different outcomes? *European Journal of Human Genetics* 29(5), 789–799. doi:10.1038/s41431-020-00799-1
- Medendorp NM, Stiggelbout AM, Aalfs CM, *et al.* (2021b) A scoping review of practice recommendations for clinicians' communication of uncertainty. *Health Expectations* 24(4), 1025–1043. doi:10.1111/hex.13255
- Medendorp N, Visser L, Hillen M, *et al.* (2017) How oncologists' communication improves (analogue) patients' recall of information. A randomized video-vignettes study. *Patient Education and Counseling* 100(7), 1338–1344. doi:10.1016/j.pec.2017.02.012
- Mishel MH (1981) The measurement of uncertainty in illness. *Nursing Research* 30(5), 258–263. doi:10.1097/00006199-198109000-00002
- Mishel MH and Clayton MF (2008) Theories of uncertainty in illness. *Middle Range Theory for Nursing* 3, 53–86.
- Mori M, Fujimori M, van Vliet LM, *et al.* (2019) Explicit prognostic disclosure to Asian women with breast cancer: A randomized, scripted video-vignette study (J-SUPPORT1601). *Cancer* 125(19), 3320–3329. doi:10.1002/cncr.32327
- Nakajima N, Hata Y, Onishi H, *et al.* (2013) The evaluation of the relationship between the level of disclosure of cancer in terminally ill patients with cancer and the quality of terminal care in these patients and their families using the Support Team Assessment Schedule. *American Journal of Hospice and Palliative Medicine* 30(4), 370–376. doi:10.1177/1049909112452466
- Nipp RD, El-Jawahri A, Fishbein JN, *et al.* (2016) The relationship between coping strategies, quality of life, and mood in patients with incurable cancer. *Cancer* 122(13), 2110–2116. doi:10.1002/cncr.30025
- Nishioka M, Okuyama T, Uchida M, *et al.* (2019) What is the appropriate communication style for family members confronting difficult surrogate decision-making in palliative care?: A randomized video vignette study in medical staff with working experiences of clinical oncology. *Japanese Journal of Clinical Oncology* 49(1), 48–56. doi:10.1093/jjco/hyy178
- Olsman E, Leget C, Onwuteaka-Philipsen B, *et al.* (2014) Should palliative care patients' hope be truthful, helpful or valuable? An interpretative synthesis of literature describing healthcare professionals' perspectives on hope of palliative care patients. *Palliative Medicine* 28(1), 59–70. doi:10.1177/0269216313482172
- Parker SM, Clayton JM, Hancock K, *et al.* (2007) A systematic review of prognostic/end-of-life communication with adults in the advanced stages of a life-limiting illness: Patient/caregiver preferences for the content, style, and timing of information. *Journal of Pain and Symptom Management* 34(1), 81–93. doi:10.1016/j.jpainsymman.2006.09.035
- Rains SA, Tukachinsky R (2015) Information seeking in uncertainty management theory: Exposure to information about medical uncertainty and information-processing orientation as predictors of uncertainty management success. *Journal of Health Communication* 20(11), 1275–1286. doi:10.1080/10810730.2015.1018641
- Reis-Dennis S, Gerrity MS and Geller G (2021) Tolerance for uncertainty and professional development: A normative analysis. *Journal of General Internal Medicine* 36(8), 1–6. doi:10.1007/s11606-020-06538-y
- Scheier MF, Carver CS, Bridges MW, *et al.* (1994) Distinguishing optimism from neuroticism (and trait anxiety, self-mastery, and self-esteem): A reevaluation of the Life Orientation Test. *Journal of Personality and Social Psychology* 67(6), 1063–1078. doi:10.1037/0022-3514.67.6.1063
- Shilling V, Starkings R, Jenkins V, *et al.* (2017) The pervasive nature of uncertainty—A qualitative study of patients with advanced cancer and their informal caregivers. *Journal of Cancer Survivorship* 11(5), 590–603. doi:10.1007/s11764-017-0628-x
- Shin JA, El-Jawahri A, Parkes A, *et al.* (2016) Quality of life, mood, and prognostic understanding in patients with metastatic breast cancer. *Journal of Palliative Medicine* 19(8), 863–869. doi:10.1089/jpm.2016.0027

- Sisk BA and Mack JW** (2018) How should we enhance the process and purpose of prognostic communication in oncology? *AMA Journal of Ethics* **20**(8), 757–765. doi:10.1001/amajethics.2018.757
- Stiggelbout AM, Dehaes JCJM, Kiebert GM, et al.** (1996) Tradeoffs between quality and quantity of life: Development of the QQ questionnaire for cancer patient attitudes. *Medical Decision Making* **16**(2), 184–192. doi:10.1177/0272989X9601600211
- Street RL, Elwyn G and Epstein RM** (2012) Patient preferences and healthcare outcomes: An ecological perspective. *Expert Review of Pharmacoeconomics and Outcomes Research* **12**(2), 167–180. doi:10.1586/erp.12.3
- Surbone A** (2008) Cultural aspects of communication in cancer care. *Supportive Care in Cancer* **16**, 235–240. doi:10.1007/s00520-007-0366-0
- Tabachnick BG, Fidell LS and Ullman JB** (2007) *Using Multivariate Statistics*. Boston, MA: Pearson.
- Taha S, Matheson K, Cronin T, et al.** (2014) Intolerance of uncertainty, appraisals, coping, and anxiety: The case of the 2009 H1N1 pandemic. *British Journal of Health Psychology* **19**(3), 592–605. doi:10.1111/bjhp.12058
- Ter Veer E, Haj Mohammad N, van Valkenhoef G, et al.** (2016) The efficacy and safety of first-line chemotherapy in advanced esophagogastric cancer: A network meta-analysis. *Journal of the National Cancer Institute* **108**(10), djw166. doi:10.1093/jnci/djw166
- van der Velden NC, Meijers MC, Han PK, et al.** (2020) The effect of prognostic communication on patient outcomes in palliative cancer care: A systematic review. *Current Treatment Options in Oncology* **21**(5), 1–38. doi:10.1007/s11864-020-00742-y
- van der Velden NC, Smets EM, van Vliet LM, et al.** (2024) Effects of prognostic communication strategies on prognostic perceptions, treatment decisions and end-of-life anticipation in advanced cancer: An experimental study among analogue patients. *Journal of Pain and Symptom Management*, In Press. doi:10.1016/j.jpainsymman.2024.02.563
- van der Velden NC, van der Kleij M, Lehmann V, et al.** (2021) Communication about prognosis during patient-initiated second opinion consultations in advanced cancer care: An observational qualitative analysis. *International Journal of Environmental Research & Public Health* **18**(11), 5694. doi:10.3390/ijerph18115694
- van der Velden NC, van Laarhoven HWM, Burgers SA, et al.** (2022b) Characteristics of patients with advanced cancer preferring not to know prognosis: A multicenter survey study. *BMC Cancer* **22**(1), 941. doi:10.1186/s12885-022-09911-8
- van der Velden NC, van Laarhoven HW, Nieuwkerk PT, et al.** (2022a) Attitudes toward striving for quality and length of life among patients with advanced cancer and a poor prognosis. *JCO Oncology Practice* **18**(11), e1818–e1830. doi:10.1200/OP.22.00185
- van Osch M, Sep M, van Vliet LM, et al.** (2014) Reducing patients' anxiety and uncertainty, and improving recall in bad news consultations. *Health Psychology* **33**(11), 1382. doi:10.1037/hea0000097
- van Osch M, van Dulmen S, van Vliet LM, et al.** (2017) Specifying the effects of physician's communication on patients' outcomes: A randomised controlled trial. *Patient Education and Counseling* **100**(8), 1482–1489. doi:10.1016/j.pec.2017.03.009
- van Vliet LM, Francke A, Tomson S, et al.** (2013a) When cure is no option: How explicit and hopeful can information be given? A qualitative study in breast cancer. *Patient Education and Counseling* **90**(3), 315–322. doi:10.1016/j.pec.2011.03.021
- van Vliet LM, Hillen MA, van der Wall E, et al.** (2013b) How to create and administer scripted video-vignettes in an experimental study on disclosure of a palliative breast cancer diagnosis. *Patient Education and Counseling* **91**(1), 56–64. doi:10.1016/j.pec.2012.10.017
- van Vliet LM, van der Wall E, Albada A, et al.** (2012) The validity of using analogue patients in practitioner–patient communication research: Systematic review and meta-analysis. *Journal of General Internal Medicine* **27**(11), 1528–1543. doi:10.1007/s11606-012-2111-8
- van Vliet LM, van der Wall E, Plum NM, et al.** (2013c) Explicit prognostic information and reassurance about nonabandonment when entering palliative breast cancer care: Findings from a scripted video-vignette study. *Journal of Clinical Oncology* **31**(26), 3242–3249. doi:10.1200/JCO.2012.45.5865
- Visser LN, Bol N, Hillen MA, et al.** (2018) Studying medical communication with video vignettes: A randomized study on how variations in video-vignette introduction format and camera focus influence analogue patients' engagement. *BMC Medical Research Methodology* **18**, 1–12. doi:10.1186/s12874-018-0472-3
- Visser LN, Hillen MA, Verdam MGE, et al.** (2016a) Assessing engagement while viewing video vignettes; validation of the Video Engagement Scale (VES). *Patient Education and Counseling* **99**(2), 227–235. doi:10.1016/j.pec.2015.08.029
- Visser LN, Tollenaar MS, Bosch JA, et al.** (2016b) Analogue patients' self-reported engagement and psychophysiological arousal in a video-vignettes design: Patients versus disease-naïve individuals. *Patient Education and Counseling* **99**(10), 1724–1732. doi:10.1016/j.pec.2016.04.012
- Visser LN, Tollenaar MS, Bosch JA, et al.** (2017) Are psychophysiological arousal and self-reported emotional stress during an oncological consultation related to memory of medical information? An experimental study. *Stress* **20**(1), 103–111. doi:10.1080/10253890.2017.1286323
- Visser LN, Tollenaar MS, van Doornen LJ, et al.** (2019) Does silence speak louder than words? The impact of oncologists' emotion-oriented communication on analogue patients' information recall and emotional stress. *Patient Education and Counseling* **102**(1), 43–52. doi:10.1016/j.pec.2018.08.032
- Visser LN, van der Velden NC, Smets EM, et al.** (2022) Methodological choices in experimental research on medical communication using vignettes: The impact of gender congruence and vignette modality. *Patient Education and Counseling* **105**(6), 1634–1641. doi:10.1016/j.pec.2021.10.015
- Watson D, Clark LA and Tellegen A** (1988a) Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology* **54**(6), 1063–1070. doi:10.1037/0022-3514.54.6.1063
- Watson M, Greer S, Young J, et al.** (1988b) Development of a questionnaire measure of adjustment to cancer: The MAC scale. *Psychological Medicine* **18**(1), 203–209. doi:10.1017/S0033291700002026
- Watson M, Law M, Dossantos M, et al.** (1994) The Mini-MAC: Further development of the mental adjustment to cancer scale. *Journal of Psychosocial Oncology* **12**(3), 33–46. doi:10.1300/J077V12N03\_03
- Westendorp J, Evers AW, Stouthard JM, et al.** (2022) Mind your words: Oncologists' communication that potentially harms patients with advanced cancer: A survey on patient perspectives. *Cancer* **128**(5), 1133–1140. doi:10.1002/cncr.34018