

INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, <http://www.editorialmanager.com/psm/>. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

1. S.I. units should be used throughout in text, figures and tables.
2. Authors should spell out in full any abbreviations used in their manuscripts.
3. Foreign quotations and phrases should be followed by a translation.
4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* **286**, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* **32**, 173–179.

Cleckley HJ (1941). *The Mask of Sanity*, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank.org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq_tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. **Line artwork:** Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; **Combination artwork (line/tone):** Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; **Black and white halftone artwork:** Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; **Colour halftone artwork:** Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. *Tables* Tables should be numbered consecutively in the text in Arabic numerals and each typed on a separate sheet after the References section. Titles should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

REVIEW ARTICLES

Attention to threat in posttraumatic stress disorder as indexed by eye-tracking indices: a systematic review
Lazarov A, Suarez-Jimenez B, Tamman A, Falzon L, Zhu X, Edmondson DE & Neria Y 705

Longitudinal outcomes of patients with pseudodementia: a systematic review
Connors MH, Quinto L & Brodaty H 727

Neurocognitive and neuroimaging correlates of obesity and components of metabolic syndrome in bipolar disorder: a systematic review
Bora E, McIntyre RS & Ozerdem A 738

COMMENTARY

A critical analysis of recent data on the long-term outcome of antipsychotic treatment
Moncrieff J & Steingard S 750

ORIGINAL ARTICLES

Cortical structures and their clinical correlates in antipsychotic-naïve schizophrenia patients before and after 6 weeks of dopamine D_{2/3} receptor antagonist treatment
Jessen K, Rostrup E, Mandl RCW, Nielsen MØ, Bak N, Fagerlund B, Glenthøj BY & Ebdrup BH 754

Dimensional analysis of depressive, anxious and somatic symptoms presented by primary care patients and their relationship with ICD-11 PHC proposed diagnoses
Ziebold C, Goldberg DP, Reed GM, Minhas F, Razzaque B, Fortes S, Robles R, Lam TP, Bobes J, Iglesias C, Cogo-Moreira H, García JA & Mari JJ 764

Effect of discontinuation v. maintenance of antipsychotic medication on relapse rates in patients with remitted/stable first-episode psychosis: a meta-analysis
Kishi T, Ikuta T, Matsui Y, Inada K, Matsuda Y, Mishima K & Iwata N 772

Learning to trust: social feedback normalizes trust behavior in first-episode psychosis and clinical high risk
Lemmers-Jansen ILJ, Fett A-KJ, Hanssen E, Veltman DJ & Krabbendam L 780

Posttraumatic psychopathology and the pace of the epigenetic clock: a longitudinal investigation
Wolf EJ, Logue MW, Morrison FG, Wilcox ES, Stone A, Schichman SA, McGlinchey RE, Milberg WP & Miller MW 791

Modulation of orbitofrontal-striatal reward activity by dopaminergic functional polymorphisms contributes to a predisposition to alcohol misuse in early adolescence
Baker TE, Castellanos-Ryan N, Schumann G, Cattrell A, Flor H, Nees F, Banaschewski T, Bokde A, Whelan R, Buechel C, Bromberg U, Orfanos DP, Gallinat J, Garavan H, Heinz A, Walter H, Brühl R, Gowland P, Paus T, Poustka L, Martinot J-L, Lemaître H, Artiges E, Martinot M-LP, Smolka MN, Conrod P & the IMAGEN consortium 801

The symptomatology of psychological trauma in the aftermath of war (1945–1980): UK army veterans, civilians and emergency responders
Engelbrecht A, Burdett H, Silva MJ, Bhui K & Jones E 811

Patterns of diagnostic transition in eating disorders: a longitudinal population study in Sweden
Schaumberg K, Jangmo A, Thornton LM, Birgegård A, Almqvist C, Norring C, Larsson H & Bulik CM 819

The effectiveness and cost-effectiveness of the Incredible Years® Teacher Classroom Management programme in primary school children: results of the STARS cluster randomised controlled trial
Ford T, Hayes R, Byford S, Edwards V, Fletcher M, Logan S, Norwich B, Pritchard W, Allen K, Allwood M, Gangui P, Grimes K, Hansford L, Longdon B, Norman S, Price A & Ukoumunne OC 828

Associations of trauma exposure and post-traumatic stress disorder with the activity of the renin-angiotensin-aldosterone-system in the general population
Terock J, Hannemann A, Janowitz D, Freyberger HJ, Felix SB, Dörr M, Nauck M, Völzke H & Grabe HJ 843

More than just statics: temporal dynamics of intrinsic brain activity predicts the suicidal ideation in depressed patients
Li J, Duan X, Cui Q, Chen H & Liao W 852

Prolonged grief disorder: clinical utility of ICD-11 diagnostic guidelines
Mauro C, Reynolds III CF, Maercker A, Skritskaya N, Simon N, Zisook S, Lebowitz B, Cozza SJ & Shear MK 861

Associations between P3a and P3b amplitudes and cognition in antipsychotic-naïve first-episode schizophrenia patients
Kruiper C, Fagerlund B, Nielsen MØ, Düring S, Jensen MH, Ebdrup BH, Glenthøj BY & Oranje B 868

Correspondence 876, 879